

10-10-00

A

10/06/00
jc943 U.S. PTO

Patent Application Transmittal
(only for new nonprovisional applications
under 37 C.F.R. 1.53(b))
Correspondence Address:
FROMMER LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151
Tel. (212) 588-0800 Fax: (212) 588-0500

jc925 U.S. PTO
09/680228
10/06/00

Re: 454313-2340.2

TO: ASSISTANT COMMISSIONER FOR PATENTS
Box PATENT APPLICATION
Washington, D.C. 20231

Sir:

With reference to the filing in the United States Patent and Trademark Office of an application for patent in the name of:

ELLIS ET AL.

(information on attached UNSIGNED Declaration & Power of Attorney)

entitled: PREVENTION OF MYOCARDITIS, ABORTION AND INTRAUTERINE
INFECTION ASSOCIATED WITH PORCINE CIRCOVIRUS-2

☐ This is an application of a small entity under 37 CFR 1.9(f).

☐ Small Entity Verified Statement is enclosed (unsigned)

The following are enclosed:

- ☒ Specification (51 pages, plus 1 page of abstract on page 60.)
- ☒ (8) Sheet(s) of Drawings (FIGURES 1-8)
- ☒ 80 Claims (pp. 52-59)
- ☒ Oath or Declaration and Power of Attorney (unsigned)
- ☒ Filing Fee for 3 independent claims and 81 claims in total of: \$1,458.00 for a total of \$2,148.00
- ☒ An executed Declaration and Power of Attorney will be filed upon receipt of a Notice To File Missing Parts. Kindly accord the application an October 6, 2000 filing date.

Respectfully submitted,
FROMMER LAWRENCE & HAUG LLP
Attorneys for Applicants

By: Thomas J. Kowalski
Thomas J. Kowalski, Reg. No. 32,147

EXPRESS MAIL

Mailing Label Number: EL588272455US

Date of Deposit: October 6, 2000

I hereby certify that this paper or fee is being deposited with the
United States Postal Service "Express Mail Post Office to
Addressee" Service under 37 CFR 1.10 on the date indicated
above and is addressed to: Assistant Commissioner for Patents,
Washington, D.C. 20231.

Edward Nay
(Typed or printed name of person mailing paper or fee)

Edward Nay
(Signature of person mailing paper or fee)

TITLE OF THE INVENTION

***PREVENTION OF MYOCARDITIS, ABORTION
AND INTRAUTERINE INFECTION ASSOCIATED
WITH PORCINE CIRCOVIRUS-2***

RELATED APPLICATIONS

This application is a Continuation-in-Part of U.S. Application Serial No. 09/583,350, filed May 31, 2000, which claims priority from and is based upon U.S. application Serial No. 60/151,564, filed August 31, 1999 and, this application claims priority from PCT/FR0002392, filed August 28, 2000.

Reference is also made to: WO-A-0 1409, published from PCT/EP99/04698, filed June 28, 1999 and U.S. application Serial No. 09/347,594, filed July 1, 1999, both claiming priority from French application No. 98 08777, filed July 6, 1998; U.S. application Serial No. 09/161,092, filed September 25, 1998 as a continuation-in-part of U.S. application Serial No. 09/082,558, filed May 21, 1998, claiming priority from French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively; WO-A-99 18214; and the U.S. applications of Audonnet et al. and Bublot et al., Serial Nos. 60/138,352 and 60/138,478, respectively, both filed June 10, 1999, and Serial Nos. 09/586,535 and 09/583,545, filed May 31 and June 1, 2000, respectively ("DNA VACCINE-PCV", and "PORCINE CIRCOVIRUS RECOMBINANT POXVIRUS VACCINE"). Reference is additionally made to each of the documents cited in the text and in the record or prosecution of each of the aforementioned U.S. and French applications, including without limitation WO 98/03658, published January 29, 1998 from PCT/FR97/01313, filed July 15, 1997 and designating the U.S. and claiming priority from French application 96 09338, filed July 19, 1996 (the U.S. continuation-in-part of PCT/FR97/01313 being U.S. application Serial No. 09/232,468, filed January 15, 1999 for "POLYNUCLEOTIDE VACCINE FORMULA AGAINST PORCINE REPRODUCTIVE AND RESPIRATORY PATHOLOGIES"). Mention is also made of PCT WO99/29717. Each of the aforementioned U.S., PCT and French applications (including parenthetically), and each document cited in the text and the record or prosecution of each of the aforementioned U.S., PCT and French applications (including parenthetically) ("application cited documents") and each document cited or referenced in each of the application cited documents, is hereby incorporated herein by reference; and, technology in each of the aforementioned U.S., PCT and French applications (including parenthetically), and each document cited in the text and

the record or prosecution of each of the aforementioned U.S., PCT and French applications (including parenthetically) can be used in the practice of this invention.

FIELD OF THE INVENTION

The invention relates to methods and/or compositions for the prevention and/or treatment of PCV-2-caused myocarditis, and/or abortion and/or intrauterine infection, as well as pathologic sequelae including but not limited to post-weaning multisystemic wasting syndrome; and, to methods for preparing such compositions and kits for preparing such compositions or for performing such methods, *inter alia*.

Various documents are cited in this text. Citations in the text can be by way of a citation to a document in the reference list, e.g., by way of an author(s) and document year citation to a document listed in the reference list, or by full citation in the text to a document that may or may not also be listed in the reference list.

There is no admission that any of the various documents cited in this text are prior art as to the present invention. Any document having as an author or inventor person or persons named as an inventor herein is a document that is not by another as to the inventive entity herein. All documents cited in this text ("herein cited documents") and all documents cited or referenced in herein cited documents are hereby incorporated herein by reference.

BACKGROUND OF THE INVENTION

Porcine circovirus-2 (PCV-2) was recently identified as an agent that has been consistently associated with post-weaning multisystemic wasting syndrome (PMWS) in swine populations in several parts of the world (Allan et al. 1998; Ellis et al., 1998). Isolates of PCV-2 obtained from infected pigs in several countries are virtually identical genetically, and are distinctly different from the PCV (CCL33, PCV-1) that was originally identified in the 1970's as a noncytopathic contaminant of porcine kidney (PK/15) cell line (Meehan et al. 1998; Tischer et al. 1974). Pigs with naturally acquired or experimentally induced PCV-2 infections present with progressive weight loss, tachypnea, dyspnea, and jaundice (Allan et al. 1998; Allan et al. 1999; Ellis et al. 1998; Ellis et al. 1999). Gross pathologic findings that have been directly associated with PCV-2 antigen include, lymphadenopathy, interstitial pneumonia, hepatitis and nephritis (Allan et al. 1998; Allan et al. 1999; Ellis et al. 1998; Ellis et al. 1999). PCV-2 has not heretofore been directly linked to abortion or lesions in fetal pigs. Thus, heretofore, it has not

been proposed to address the issue of PCV-2-caused myocarditis, and/or abortion and/or intrauterine infection.

OBJECTS AND SUMMARY OF THE INVENTION

It has surprisingly been found that PCV-2 is a causative agent of myocarditis, abortion and intrauterine infection, as well as post-weaning multisystemic wasting syndrome.

By definition, a PCV-2 immunogen is intended to encompass live attenuated or inactivated PCV-2, or subunit(s) from PCV-2 obtained by *in vitro* expression or by extraction, or fragment(s) comprising at least one epitope of interest which can be obtained by chemical synthesis or by *in vitro* recombinant expression, as well as recombinant vector(s) comprising and expressing *in vivo* sequence(s) or fragment(s) or epitope(s) of PCV-2 genome as herein disclosed or as in documents cited or referenced herein.

A similar definition applies for an immunogen of another porcine pathogen as disclosed herein.

Thus, an object of the invention can be to provide methods and/or compositions for the prevention and/or treatment of PCV-2-caused myocarditis, and/or abortion and/or intrauterine infection, as well as post-weaning multisystemic wasting syndrome and/or pathologic sequelae including but not limited to post-weaning multisystemic wasting syndrome; and, methods for formulating such compositions and uses of a PCV-2 immunogen (which compositions can also include a porcine parvovirus (PPV) immunogen, wherein when recombinant vector expression is used, the vector can co-express both the PPV and the PCV-2 immunogens, *inter alia*) for formulating such compositions.

Another object of the invention is the isolation and characterisation of new PCV-2 strains identified 1103 (1103/1 P.2) and 1121 (1121/1 P.1), and their uses to produce immunogens, as well as antigens and antibodies for diagnostics, in relation with PCV-2-caused myocarditis, and/or abortion and/or intrauterine infection, as well as post-weaning multisystemic wasting syndrome and/or pathologic sequelae associated therewith.

The invention provides also for inoculation of female pigs (e.g., sows, gilts) with a composition comprising a (at least one) PCV-2 immunogen (which composition can also include an immunogen from porcine parvovirus) prior to breeding; and/or prior to serving, and/or during gestation (or pregnancy); and/or prior to the perinatal period or farrowing; and/or repeatedly over a lifetime, to prevent myocarditis and/or abortion and/or intrauterine infection associated with

PCV-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2; or, to elicit an immunogenic or protective response against PCV-2 and thereby prevent post-weaning multisystemic wasting syndrome and/or myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2 and/or other pathologic sequelae associated with PCV-2.

Advantageously, at least one inoculation is done before serving. It is also advantageously followed by an inoculation to be performed during gestation, e.g., at about mid-gestation (at about 6-8 weeks of gestation) and/or at the end of gestation (at about 11-13 weeks of gestation). Thus, an advantageous regimen is an inoculation before serving and a booster inoculation during gestation. Thereafter, there can be reinoculation before each serving and/or during gestation at about mid-gestation (at about 6-8 weeks of gestation) and/or at the end of gestation (at about 11-13 weeks of gestation). Preferably, reinoculation can be during gestation only.

In another preferred embodiment, piglets, such as piglets from vaccinated females (e.g., inoculated as herein discussed), are inoculated within the first weeks of life, e.g., inoculation at one and/or two and/or three and/or four and/or five weeks of life. More preferably, piglets are first inoculated within the first week of life or within the third week of life (e.g., at the time of weaning). Even more advantageous, such piglets are then boosted two (2) to four (4) weeks later (after being first inoculated). Thus, both offspring, as well as female pig (e.g., sow, gilt) can be administered compositions of the invention and/or can be the subject of performance of methods of the invention.

Thus, the invention also comprehends immunogenic or vaccine compositions for preventing or treating myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2. An immunogenic (or immunological) composition elicits an immunological response - local or systemic. A vaccine composition elicits a local or systemic protective response. The terms "immunological composition" and "immunogenic composition" include a "vaccine composition" (as the two former terms can be protective compositions). The composition can comprise a PCV-2 immunogen (which composition can also include a PPV immunogen).

And, the invention further comprehends uses of a PCV-2 immunogen (which composition can also include a PPV immunogen) to formulate an immunogenic or vaccine

Further still, the invention comprehends an immunogenic or vaccine composition for the prevention and/or treatment of PCV-2-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome comprising a pharmaceutically or veterinarily acceptable carrier and/ or vehicle and/or excipient and/or adjuvant, and a PCV-2 immunogen

The composition can additionally include at least one immunogen from at least one additional pig pathogen, e.g.: Porcine Reproductive and Respiratory Syndrome (PRRS), *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, *Bordetella bronchiseptica*, *Pasteurella multocida*, *Erysipelothrix rhusiopathiae*, Pseudorabies, Hog cholera, Swine Influenza, and Porcine Parvovirus (PPV). Thus, vector-based compositions can include at least one immunogen from at least one additional pig pathogen, such as a vector expressing a sequence from this pathogen, wherein the vector can also be the vector expressing the PCV-2 immunogen. The vector expressing a PCV-2 sequence can comprise a PCV-2 sequence or fragment thereof as herein disclosed or as in documents cited or referenced herein; and the invention comprehends such nucleic acid molecules, vectors containing them, compositions comprising such nucleic acid molecules or vector expression products from such nucleic acid molecules, compositions comprising such expression products, probes or primers for such nucleic acid molecules, and methods for making and using any or all of the foregoing.

The vector can comprise a DNA vector plasmid, a bacteria such as an *E. coli*, a virus such as baculovirus, a herpesvirus including pig herpes viruses, including Aujeszky's disease virus, an adenovirus including a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, a racoonpox and a swinepox virus, and the like. The vector-based compositions can comprise a vector that contains and expresses an ORF selected from the group consisting of ORFs 1 to 13, such as an ORF selected from ORFs 4, 7, 10 and 13; preferably ORFs 4 and/or 13, of a PCV-2, advantageously of any one of the PCV-2 strains identified herein. And, the immunogen in compositions (either PCV-2 and/or from another pig pathogen) can be recombinantly produced. The word plasmid is intended to include any DNA transcription unit in the form of a polynucleotide sequence comprising the PCV sequence to be expressed.

Advantageously, the plasmid includes elements necessary for its expression; for instance, expression *in vivo*. The circular plasmid form, supercoiled or otherwise, is advantageous; and, the linear form is also included within the scope of the invention. The plasmid immunogenic or vaccine composition can be administered by way of a gene gun, intradermally via an needleless injector, subcutaneously or intramuscularly, or by mucosal route, or by any other means that allows for expression *in vivo*, and advantageously an immunogenic or protective response.

It is noted that the expression product generated by vectors or recombinants in this invention optionally can also be isolated and/or purified from infected or transfected cells; for instance, to prepare compositions for administration to pigs; however, in certain instances, it may be advantageous not to isolate and/or purify an expression product from a cell; for instance, when the cell or portions thereof enhance the immunogenic effect of the polypeptide. And, techniques for protein purification and/or isolation from this disclosure and documents cited herein, *inter alia*, and thus within the ambit of the skilled artisan, can be used, without undue experimentation, to purify and/or isolate recombinant or vector expression products and/or subunits of PCV-2 and/or other pig pathogens, in the practice of the invention, and such techniques, in general, can include: precipitation by taking advantage of the solubility of the protein of interest at varying salt concentrations, precipitation with organic solvents, polymers and other materials, affinity precipitation and selective denaturation; column chromatography, including high performance liquid chromatography (HPLC), ion-exchange, affinity, immunoaffinity or dye-ligand chromatography; immunoprecipitation, gel filtration, electrophoretic methods, ultrafiltration and isoelectric focusing, and their combinations, *inter alia*.

The invention further envisages methods for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising inducing an immunogenic or protective response against PCV-2 in a pig comprising administering to the pig an aforementioned or herein disclosed composition.

Thus, the invention comprehends a method for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising inducing an immunogenic or protective response against PCV-2 in a pig

comprising administering to the pig a composition comprising a pharmaceutically or veterinarily acceptable carrier or excipient or vehicle, with preferably an adjuvant, and an active agent comprising a PCV-2 immunogen. The method can be for the prevention of PCV-2-caused myocarditis and/or abortion and/or intrauterine infection comprising administering a composition comprising a pharmaceutically or veterinarily acceptable carrier and a PCV-2 immunogen. The PCV-2 immunogen can be an attenuated live whole PCV-2 or inactivated PCV-2. The method can involve a composition that is a subunit immunogenic, or vaccine composition. The method can involve the composition additionally including at least one immunogen from at least one additional pig pathogen, including a vector expressing such an immunogen or epitope; e.g., the at least one additional pig pathogen can be selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, *Pseudorabies*, Hog cholera, *Bordetella bronchiseptica*, *Pasteurella multocida*, *Erysipelothrix rhusiopathiae*, Swine Influenza, and PPV and combinations thereof. The method can involve a vector that is a DNA vector plasmid, a bacteria such as an *E. coli*, a virus such as baculovirus, a herpesvirus including Aujeszky's disease virus, an adenovirus including a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, and a swinepox virus, and the like. The method can involve a vector-based composition additionally including at least one sequence, fragment or epitope from at least one additional pig pathogen, or a vector expressing such a sequence, fragment or epitope, wherein the vector can also be the vector expressing the PCV-2 sequence, fragment or epitope. The method can involve a vector that contains and expresses an ORF selected from the group consisting of ORFs 1 to 13, e.g., an ORF selected from ORFs 4, 7, 10 and 13; preferably ORFs 4 and/or 13. The method can also involve an immunogen-based composition wherein one or more of the immunogen(s) is recombinantly produced. In this method, females and/or piglets are inoculated as described above.

In another embodiment, the invention involves a method for preparing any of the aforementioned or herein disclosed compositions comprising admixing the pharmaceutically or veterinarily acceptable carrier and the PCV-2 immunogen. The method can further include transfecting or infecting a cell or host with a recombinant vector that contains DNA encoding a PCV-2 immunogen and expresses that immunogen; and optionally purifying and/or isolating the immunogen from the cell. Similarly the method can include isolating and/or purifying a PCV-2 immunogen from PCV-2, or isolating PCV-2 from a sample.

The invention also provides a kit for preparing any of the aforementioned or herein disclosed compositions or for performing any of the aforementioned or herein disclosed methods comprising in a first container the pharmaceutically or veterinarily acceptable carrier or vehicle or excipient and in a second container the active agent comprising the PCV-2 immunogen, wherein the first and second containers are optionally packaged together, and the kit optionally includes instructions for admixture of ingredients of the composition and/or administration of the composition.

In yet another embodiment, the invention provides for administering any of the aforementioned or herein disclosed compositions to male and/or female pigs; to prevent transmission of PCV-2 and prevent or treat or control myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2. Administration is preferably done as described above.

The term “comprising” in this disclosure can mean “including” or can have the meaning commonly given to the term “comprising” in U.S. Patent Law.

Other aspects of the invention are described in or are obvious from (and within the ambit of the invention) the following disclosure.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 depicts SEQ ID No. 1, the PCV DNA sequence of the genome of the Imp. 1011-48121 strain.

Figure 2 depicts SEQ ID No. 2, the DNA sequence of the genome of the Imp. 1011-48285 strain.

Figure 3 depicts SEQ ID No. 3, the DNA sequence of the genome of the Imp. 999 strain.

Figure 4 depicts SEQ ID No. 4, the DNA sequence of the genome of the Imp. 1010 strain.

Figure 5 depicts SEQ ID No. 5, the DNA sequence of the genome of the PK/15 strain.

Figure 6 depicts SEQ ID No. 6, the DNA sequence of the genome of the Imp. 999 strain as defined in the first filing in France on October 3, 1997.

Figure 7 depicts SEQ ID No. 7, the DNA sequence of the genome of the 1103 strain, isolated in Alberta, Canada. k means g (G) or t (T), y means c (C) or t (T). These variations of sequence were observed in the viral population.

Figure 8 depicts SEQ ID No. 8, the DNA sequence of the genome of the 1121 strain, isolated in Saskatoon, Canada.

DETAILED DESCRIPTION

Porcine circovirus-2 (PCV-2) is an agent associated with post-weaning multisystemic wasting syndrome (PMWS) in swine populations. As shown in Examples 1 and 2, the potential spectrum of disease associated with PCV-2 is expanded by evidence of vertical transmission and associated reproductive failure.

In particular, Example 1 shows that PCV-2 was isolated from a litter of aborted piglets from a farm experiencing late term abortions and stillbirths. Severe, diffuse myocarditis was present in one piglet associated with extensive immunohistochemical staining for PCV-2 antigen. Variable amounts of PCV-2 antigen were also present in liver, lung and kidney of multiple fetuses. The presence of other agents that have been associated with fetal lesions and abortion in swine including porcine parvovirus, porcine reproductive respiratory syndrome virus, encephalomyocarditis virus and enterovirus could not be established.

More in particular, Example 2 shows that tissues obtained from 30 high health herds over a four-year period, and tested in routine cases of abortion or reproductive failure, were positive for PCV-2 in two submissions involving several stillborn piglets and non-viable neonates presenting with severe diffuse myocarditis, cardiac hypertrophy and evidence of chronic passive congestion. The two positive submissions were from the same farm, but occurred at two different times. The presence of PCV-2 in the hearts and other tissues of affected piglets was confirmed by immunohistochemistry and virus isolation. Failure to detect porcine circoviruses in cases of reproductive failure prior to 1999 in areas of endemic infections supports the view that reproductive disease is a new clinical manifestation of PCV-2 infection, and further suggests that sexual, as well as vertical, modes of transmission are responsible for viral dissemination in the pig population.

Accordingly, inoculation of pigs, e.g., female pigs, such as sows or gilts, with a composition including at least one PCV-2 immunogen (e.g. from at least one strain chosen among strains Imp1008, Imp1010, Imp999, Imp1011-48285, Imp1011-48121, 1103 and 1121) (which composition can also include at least one immunogen from at least one other porcine pathogen such as at least one porcine parvovirus, wherein when a vector is used the vector can co-express both the PCV-2 immunogen(s) and the at least one immunogen of the at least one

other porcine pathogen, e.g., PPV immunogen(s), *inter alia*), in a schedule of immunization as described above, can prevent myocarditis and/or abortion and/or intrauterine infection associated with PCV-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2.

Thus, the invention involves methods and compositions using PCV-2 immunogen for preventing myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2. In particular, immunogen from strain 1103 and/or strain 1121 is useful for methods and compositions using PCV-2 immunogen for preventing myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2

The PCV-2 immunogen can be any PCV-2 immunogen including any PCV-2-expressing vector identified in any herein cited document (or any document cited in herein cited documents) including any or all of: U.S. application Serial No. 09/347,594, filed July 1, 1999; French application No. 98 08777, filed July 6, 1998; U.S. application Serial No. 09/161,092, filed September 25, 1998; U.S. application Serial No. 09/082,558, filed May 21, 1998; French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively; WO-A-99 18214; the U.S. applications of Audonnet et al. and Bublot et al., Serial Nos. 60/138,352 and 60/138,478, respectively, both filed June 10, 1999 ("DNA VACCINE-PCV", and "PORCINE CIRCOVIRUS RECOMBINANT POXVIRUS VACCINE", respectively); and WO99/29717 (all of which and documents cited therein and in the prosecution thereof being hereby incorporated herein by reference). Thus, the immunogen from PCV-2 including a vector expressing such an immunogen can be prepared in accordance with herein cited documents (or documents cited in herein cited documents).

The composition comprising the PCV-2 immunogen employed in the practice of this invention can be as in any herein cited document (or any document cited in herein cited documents) including any or all of: U.S. application Serial No. 09/347,594, filed July 1, 1999; French application No. 98 08777, filed July 6, 1998; U.S. application Serial No. 09/161,092, filed September 25, 1998; U.S. application Serial No. 09/082,558, filed May 21, 1998; French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively; WO-A-99 18214; the U.S. applications of Audonnet et al. and Bublot et al., Serial Nos. 60/138,352 and 60/138,478, respectively, both filed June 10, 1999, and

Serial Nos. 09/586,535 and 09/583,545, filed May 31 and June 1, 2000, respectively ("DNA VACCINE-PCV", and "PORCINE CIRCOVIRUS RECOMBINANT POXVIRUS VACCINE", respectively); and WO99/29717 (all of which and documents cited therein and in the prosecution thereof being hereby incorporated herein by reference). Thus, the composition comprising the PCV-2 immunogen including the vector expressing PCV-2 immunogen can be prepared as in herein cited documents.

The at least one immunogen from at least one other porcine pathogen can be as described in any of the aforementioned or herein cited patent or literature publications (or documents cited therein), or as used in known porcine vaccines or immunogenic compositions, or as in WO 98/03658, published January 29, 1998 from PCT/FR97/01313, filed July 15, 1997; or French application 96 09338, filed July 19, 1996; or U.S. application Serial No. 09/232,468, filed January 15, 1999 for ("POLYNUCLEOTIDE VACCINE FORMULA AGAINST PORCINE REPRODUCTIVE AND RESPIRATORY PATHOLOGIES").

The amount of PCV-2 immunogen in compositions employed in the invention can be as described in any of the aforementioned or herein cited patent or literature publications (or documents cited therein). And, the amount of at least one immunogen from at least one other porcine pathogen can be as described in any of the aforementioned or herein patent or literature publications (or documents cited therein), or as used in known porcine vaccines or immunogenic compositions.

Compositions for use in the invention can be prepared in accordance with standard techniques well known to those skilled in the veterinary or pharmaceutical or arts. Such compositions can be administered in dosages and by techniques well known to those skilled in the veterinary arts taking into consideration such factors as the age, sex, weight, condition and particular treatment of the pig, and the route of administration. The compositions can be administered alone, or can be co-administered or sequentially administered with other compositions of the invention (e.g., other compositions comprising a PCV-2 immunogen) or with other prophylactic or therapeutic compositions (e.g., other porcine immunogenic or vaccine compositions). Thus, the invention also provides multivalent or "cocktail" or combination compositions and methods employing them. In this regard, reference is made to U.S. Patent No. 5,843,456, incorporated herein by reference, and directed to rabies compositions and combination compositions and uses thereof.

Compositions of the invention may be used for parenteral or mucosal administration, preferably by intradermal or intramuscular routes. In particular for intradermal route, injection can be done using a needleless injector. When mucosal administration is used, it is possible to use oral, nasal, or ocular routes.

In such compositions the immunogen(s) may be in a mixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose or the like, and/or preferably with an adjuvant. The compositions can also be lyophilized or frozen. The compositions can contain auxiliary substances such as pH buffering agents, adjuvants, preservatives, polymer excipients used for mucosal routes, and the like, depending upon the route of administration and the preparation desired.


Standard texts, such as "REMINGTON'S PHARMACEUTICAL SCIENCE", 17th edition, 1985, incorporated herein by reference, may be consulted to prepare suitable preparations, without undue experimentation. Suitable dosages can also be based upon the text herein and documents cited herein.

Adjuvants are substances that enhance the immune response to immunogens. Adjuvants, can include aluminum hydroxide and aluminum phosphate, saponins e.g., Quil A, water-in-oil emulsion, oil-in-water emulsion, water-in-oil-in-water emulsion. The emulsion can be based in particular on light liquid paraffin oil (European Pharmacopea type); isoprenoid oil such as squalane or squalene; oil resulting from the oligomerization of alkenes, in particular of isobutene or decene; esters of acids or of alcohols containing a linear alkyl group, more particularly plant oils, ethyl oleate, propylene glycol di(caprylate/caprata), glyceryl tri(caprylate/caprata) or propylene glycol dioleate; esters of branched fatty acids or alcohols, in particular isostearic acid esters. The oil is used in combination with emulsifiers to form the emulsion. The emulsifiers are preferably nonionic surfactants, in particular esters of sorbitan, of mannide (e.g. anhydromannitol oleate), of glycerol, of polyglycerol, of propylene glycol and of oleic, isostearic, ricinoleic or hydroxystearic acid, which are optionally ethoxylated, and polyoxypropylene-polyoxyethylene copolymer blocks, in particular the Pluronic® products, especially L121. See Hunter et al., The Theory and Practical Application of Adjuvants (Ed. Stewart-Tull, D.E.S.). John Wiley and Sons, NY, pp51-94 (1995) and Todd et al., Vaccine 15:564-570 (1997).

A further instance of an adjuvant is a compound chosen from the polymers of acrylic or methacrylic acid and the copolymers of maleic anhydride and alkenyl derivative. Advantageous adjuvant compounds are the polymers of acrylic or methacrylic acid which are cross-linked, especially with polyalkenyl ethers of sugars or polyalcohols. These compounds are known by the term carbomer (Phameuropa Vol. 8, No. 2, June 1996). Persons skilled in the art can also refer to U.S. Patent No. 2,909,462 (incorporated herein by reference) which describes such acrylic polymers cross-linked with a polyhydroxylated compound having at least 3 hydroxyl groups, preferably not more than 8, the hydrogen atoms of at least three hydroxyls being replaced by unsaturated aliphatic radicals having at least 2 carbon atoms. The preferred radicals are those containing from 2 to 4 carbon atoms, e.g. vinyls, allyls and other ethylenically unsaturated

groups. The unsaturated radicals may themselves contain other substituents, such as methyl. The products sold under the name Carbopol® (BF Goodrich, Ohio, USA) are particularly appropriate. They are cross-linked with an allyl sucrose or with allyl pentaerythritol. Among then, there may be mentioned Carbopol® 974P, 934P and 971P. Among the copolymers of maleic anhydride and alkenyl derivative, the copolymers EMA® (Monsanto) which are copolymers of maleic anhydride and ethylene, linear or cross-linked, for example cross-linked with divinyl ether, are preferred. Reference may be made to J. Fields et al., Nature, **186** : 778-780, 4 June 1960, incorporated herein by reference.

From the point of view of their structure, the polymers of acrylic or methacrylic acid and the copolymers EMA® are preferably formed of basic units of the following formula :



in which:

R₁ and R₂, which are identical or different, represent H or CH₃;

x = 0 or 1, preferably x = 1; and

y = 1 or 2, with x + y = 2.

For the copolymers EMA®, x = 0 and y = 2. For the carbomers, x = y = 1.

The dissolution of these polymers in water leads to an acid solution that will be neutralized, preferably to physiological pH, in order to give the adjuvant solution into which the immunogenic, immunological or vaccine composition itself will be incorporated. The carboxyl groups of the polymer are then partly in COO⁻ form.

Preferably, a solution of adjuvant according to the invention, especially of carbomer, is prepared in distilled water, preferably in the presence of sodium chloride, the solution obtained being at acidic pH. This stock solution is diluted by adding it to the desired quantity (for

obtaining the desired final concentration), or a substantial part thereof, of water charged with NaCl, preferably physiological saline (NaCl 9 g/l) all at once in several portions with concomitant or subsequent neutralization (pH 7.3 to 7.4), preferably with NaOH. This solution at physiological pH will be used as it is for mixing with the vaccine, which may be especially stored in freeze-dried, liquid or frozen form.

The polymer concentration in the final vaccine composition can be 0.01% to 2% w/v, e.g., 0.06 to 1% w/v, such as 0.1 to 0.6% w/v.

From this disclosure and the knowledge in the art, the skilled artisan can select a suitable adjuvant, if desired, and the amount thereof to employ in an immunological, immunogenic or vaccine composition according to the invention, without undue experimentation.

The immunogenic or vaccine compositions according to the invention may be associated to at least one live attenuated, inactivated, or sub-unit vaccine, or recombinant vaccine (e.g. poxvirus as vector or DNA plasmid) expressing at least one immunogen or epitope of interest from at least one another pig pathogen.

Compositions in forms for various administration routes are envisioned by the invention. And again, the effective dosage and route of administration are determined by known factors, such as age, sex, weight and other screening procedures which are known and do not require undue experimentation. Dosages of each active agent can be as in herein cited documents and/or can range from one or a few to a few hundred or thousand micrograms, e.g., 1 μ g to 1mg, for a subunit immunogenic, or vaccine composition; and, 10^4 to 10^{10} TCID₅₀ advantageously 10^6 to 10^8 TCID₅₀ for an inactivated (titre before inactivation) immunogenic, or vaccine composition. For a live attenuated immunogenic or vaccine composition, the dose can be between 10^1 and 10^8 TCID₅₀ advantageously 10^3 and 10^6 TCID₅₀.

Recombinants or vectors can be administered in a suitable amount to obtain *in vivo* expression corresponding to the dosages described herein and/or in herein cited documents. For instance, suitable ranges for viral suspensions can be determined empiracally. The viral vector or recombinant in the invention can be administered to a pig or infected or transfected into cells in an amount of about at least 10^3 pfu; more preferably about 10^4 pfu to about 10^{10} pfu, e.g., about 10^5 pfu to about 10^9 pfu, for instance about 10^6 pfu to about 10^8 pfu, per dose, e.g. of about 2 ml. And, if more than one gene product is expressed by more than one recombinant, each

recombinant can be administered in these amounts; or, each recombinant can be administered such that there is, in combination, a sum of recombinants comprising these amounts.

In plasmid compositions employed in the invention, dosages can be as described in documents cited herein or as described herein. For instance, suitable quantities of each plasmid DNA in plasmid compositions can be 1 µg to 2 mg, preferably 50 µg to 1mg. Documents cited herein regarding DNA plasmid vectors may be consulted by the skilled artisan to ascertain other suitable dosages for DNA plasmid vector compositions of the invention, without undue experimentation.

However, the dosage of the composition(s), concentration of components therein and timing of administering the composition(s), which elicit a suitable immunogenic response, can be determined by methods such as by antibody titrations of sera, e.g., by ELISA and/or seroneutralization assay analysis and/or by vaccination challenge evaluation in pig. Such determinations do not require undue experimentation from the knowledge of the skilled artisan, this disclosure and the documents cited herein. And, the time for sequential administrations can be likewise ascertained with methods ascertainable from this disclosure, and the knowledge in the art, without undue experimentation.

The PCV-2 immunogen can be obtained from PCV-2 or can be obtained from *in vitro* recombinant expression of PCV-2 gene(s) or portions or epitopes thereof. Methods for making and/or using vectors (or recombinants) for expression can be by or analogous to the methods disclosed in: U.S. Patent Nos. 4,603,112, 4,769,330, 5,174,993, 5,505,941, 5,338,683, 5,494,807, 4,722,848, 5,942,235, 5,364,773, 5,762,938, 5,770,212, 5,942,235, 5,756,103, 5,766,599, 6,004,777, 5,990,091, 6,033,904, 5,869,312, 5,382,425, PCT publications WO 94/16716, WO 96/39491, WO 95/30018, Paoletti, "Applications of pox virus vectors to vaccination: An update," PNAS USA 93:11349-11353, October 1996, Moss, "Genetically engineered poxviruses for recombinant gene expression, vaccination, and safety," PNAS USA 93:11341-11348, October 1996, Smith et al., U.S. Patent No. 4,745,051 (recombinant baculovirus), Richardson, C.D. (Editor), Methods in Molecular Biology 39, "Baculovirus Expression Protocols" (1995 Humana Press Inc.), Smith et al., "Production of Human Beta Interferon in Insect Cells Infected with a Baculovirus Expression Vector," Molecular and Cellular Biology, Dec., 1983, Vol. 3, No. 12, p. 2156-2165; Pennock et al., "Strong and Regulated Expression of *Escherichia coli* B-Galactosidase in Infect Cells with a Baculovirus vector," Molecular and Cellular Biology Mar.

1984, Vol. 4, No. 3, p. 399-406; EPA 0 370 573, U.S. application Serial No. 920,197, filed October 16, 1986, EP Patent publication No. 265785, U.S. Patent No. 4,769,331 (recombinant herpesvirus), Roizman, "The function of herpes simplex virus genes: A primer for genetic engineering of novel vectors," PNAS USA 93:11307-11312, October 1996, Andreansky et al., "The application of genetically engineered herpes simplex viruses to the treatment of experimental brain tumors," PNAS USA 93:11313-11318, October 1996, Robertson et al. "Epstein-Barr virus vectors for gene delivery to B lymphocytes," PNAS USA 93:11334-11340, October 1996, Frolov et al., "Alphavirus-based expression vectors: Strategies and applications," PNAS USA 93:11371-11377, October 1996, Kitson et al., J. Virol. 65, 3068-3075, 1991; U.S. Patent Nos. 5,591,439, 5,552,143, WO 98/00166, allowed U.S. applications Serial Nos. 08/675,556, and 08/675,566 both filed July 3, 1996 (recombinant adenovirus), Grunhaus et al., 1992, "Adenovirus as cloning vectors," Seminars in Virology (Vol. 3) p. 237-52, 1993, Ballay et al. EMBO Journal, vol. 4, p. 3861-65, Graham, Tibtech 8, 85-87, April, 1990, Prevec et al., J. Gen Virol. 70, 429-434, PCT WO91/11525, Felgner et al. (1994), J. Biol. Chem. 269, 2550-2561, Science, 259:1745-49, 1993 and McClements et al., "Immunization with DNA vaccines encoding glycoprotein D or glycoprotein B, alone or in combination, induces protective immunity in animal models of herpes simplex virus-2 disease," PNAS USA 93:11414-11420, October 1996, and U.S. Patents Nos 5,591,639, 5,589,466, and 5,580,859 relating to DNA expression vectors, *inter alia*. See also WO 98/33510; Ju et al., Diabetologia, 41:736-739, 1998 (lentiviral expression system); Sanford et al., U.S. Patent No. 4,945,050; Fischbach et al. (Intracel), WO 90/01543; Robinson et al., seminars in IMMUNOLOGY, vol. 9, pp.271-283 (1997) (DNA vector systems); Szoka et al., U.S. Patent No. 4,394,448 (method of inserting DNA into living cells); McCormick et al., U.S. Patent No. 5,677,178 (use of cytopathic viruses); and U.S. Patent No. 5,928,913 (vectors for gene delivery), as well as other documents cited herein. A viral vector, for instance, selected from pig herpes viruses, such as Aujeszky's disease virus, porcine adenovirus, poxviruses, especially vaccinia virus, avipox virus, canarypox virus, and swinepox virus, as well as DNA vectors (DNA plasmids) are advantageously employed in the practice of the invention .

The expression product from the PCV-2 gene(s) or portions thereof can be useful for generating antibodies such as monoclonal or polyclonal antibodies that are useful for diagnostic

purposes. Similarly, expression product(s) from the PCV-2 gene(s) or portions thereof can be useful in diagnostic applications.

Further, one skilled in the art can determine an epitope of interest in a PCV-2 immunogen, or in an immunogen of another porcine pathogen, without undue experimentation, from the disclosure herein and the knowledge in the art; *see, e.g.*, WO 98/40500, incorporated herein by reference, regarding general information for determining an epitope of interest or an epitopic region of a protein, *inter alia*.

With particular reference to U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214 (all incorporated herein by reference), particularly advantageous immunogenic, immunological or vaccine compositions are: An immunogenic or vaccine composition, collected from a cell culture *in vitro* which has been infected with a purified preparation of PCV-2, such as a purified preparation of porcine circovirus selected from the group consisting of the preparations deposited at the ECACC, under the following references: accession No. V97100219 (strain Imp.1008) , No. V97100218 (strain Imp.1010) and accession No. V97100217 (strain Imp.999) deposited October 2, 1997, accession No. V98011608(strain Imp.1011-48285) and No. V98011609 (strain Imp.1011-48121) deposited January 16, 1998, accession No. 00012710 (strain 1103) and No. 00012709 (strain 1121) deposited February 2, 2000, or an immunogenic or vaccine composition comprised of porcine circovirus produced on, and isolated from cell culture *in vitro*, these cells having been infected with a porcine circovirus capable of being isolated from a physiological sample or from a tissue sample, especially lesions, from a pig having the PMWS syndrome, e.g., such a composition wherein the porcine circovirus is produced on, and isolated from a pig kidney cell line, for instance, produced on, and isolated from PK/15 cells free from contamination with PCV-1; or such a composition comprising or prepared from a culture extract or supernatant, collected from a cell culture *in vitro* which have been infected with a such a circovirus. Thus, porcine circovirus can be an immunogen. For instance, the vaccine or immunogenic composition can comprise the attenuated live whole immunogen (e.g., virus), advantageously, in a veterinarily or pharmaceutically acceptable vehicle or diluent and optionally an a veterinarily or pharmaceutically acceptable adjuvant, as well as, optionally, a freeze-drying stabilizer. The immunogen (e.g., virus) can be inactivated

and the vaccine or immunogenic composition can additional and/or optionally comprise, a veterinarily or pharmaceutically acceptable vehicle or diluent and optionally a veterinarily or pharmaceutically acceptable adjuvant. The vaccine or immunogenic composition can comprise PCV-2 immunogens and/or immunogens of several porcine circoviruses (including PCV-2 or several strains of PCV-2, and including PCV-1), as well as optionally additionally immunogens from another pig pathogen; e.g, PRRS, Mycoplasma hyopneumoniae, Actinobacillus pleuropneumoniae, *E. coli*, Pseudorabies, Hog cholera, Bordetella bronchiseptica, Pasteurella multocida, Erysipelothrix rhusiopathiae, Swine Influenza, PPV (*see also* U.S. application Serial No. 09/347,594, filed July 1, 1999 and French application No. 98 08777, filed July 6, 1998).

For the production of circovirus antigenic preparations, the circoviruses may be obtained after passage on cells, in particular cell lines, e.g. PK/15 cells. The culture supernatants or extracts, optionally purified by standard techniques, may be used.

In the context of attenuated PCV, the attenuation may be carried out according to the customary methods, e.g. by passage on cells, preferably by passage on pig cells, especially cell lines, such as PK/15 cells (for example from 20 to 150, especially of the order of 40 to 100, passages).

In the context of inactivated vaccine, the PCV, with the fractions which may be present, is inactivated according to techniques known to persons skilled in the art. The inactivation will be preferably carried out by the chemical route, e.g. by exposing the antigen to a chemical agent such as formaldehyde (formalin), paraformaldehyde, β -propiolactone or ethyleneimine or its derivatives, and/or by physical treatment. The preferred method of inactivation will be herein the exposure to a chemical agent and in particular to ethyleneimine or to β -propiolactone.

The immunogen in the vaccine or immunogenic composition can be expressed from a DNA fragment containing a sequence or fragment thereof (advantageously encoding at least one epitope) selected from the group consisting of the sequences designated by the references SEQ ID No: 1, SEQ ID No: 2, SEQ ID No: 3, SEQ ID No: 4, SEQ ID No: 6 (in U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214), as well as SEQ ID No: 7 and SEQ ID No: 8 (Figures 1-8). The immunogen in the vaccine or immunogenic composition can be expressed from a DNA fragment containing an ORF selected from the group

consisting of ORFs 1 to 13, such as ORFs 4, 7, 10 and 13; preferably ORFs 4 and/or 13, of a PCV-2 strain, in particular of any one of the above identified strains (as designated in U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214). Thus, the immunogen or a portion thereof, such as an epitope of interest can be obtained by *in vitro* expression thereof from a recombinant or a vector. The immunogen may be further purified and/or concentrated by the conventional methods.

The immunogen in the vaccine or immunogenic composition can be expressed *in vivo* by an expression vector comprising a DNA fragment containing a sequence or fragment thereof (advantageously encoding at least one epitope) selected from the group consisting of the sequences designated by the references SEQ ID No: 1, SEQ ID No: 2, SEQ ID No: 3, SEQ ID No: 4, SEQ ID No: 6 (in U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214), as well as SEQ ID No: 7 and SEQ ID No: 8 (Figures 1-8). Similarly, the immunogen in the vaccine, immunogenic or immunological composition can be expressed *in vivo* by an expression vector comprising a DNA fragment containing an ORF selected from the group consisting of ORFs 1 to 13, such as ORFs 4, 7, 10 and 13; preferably ORFs 4 and/or 13, of a PCV-2 strain, in particular of any one of the above identified strains (as designated in U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214). That is, the vaccine or immunogenic composition can comprise an expression vector that expresses the immunogen or a portion thereof, e.g., an epitope of interest, *in vivo*.

The expression vector can be any suitable vector such as a vector selected from DNA plasmids, bacteria such as *E. coli*, viruses such as baculovirus, herpesvirus such as Aujeszky's disease virus, adenovirus including porcine adenovirus, poxviruses, especially vaccinia virus, avipox virus, canarypox virus, and swinepox virus, *inter alia* (See also the U.S. applications of Audonnet et al. and Bublot et al., Serial Nos. 60/138,352 and 60/138,478, respectively, both filed

June 10, 1999 ("DNA VACCINE-PCV", and "PORCINE CIRCOVIRUS RECOMBINANT POXVIRUS VACCINE", respectively).

Accordingly, the invention also comprehends nucleic acid molecules and vectors containing them, as well as expression products therefrom, compositions comprising such nucleic acid molecules and/or vectors and/or expression products, as well as methods for making and using any or all of these embodiments. The invention especially encompasses herein disclosed nucleic acid molecules, nucleic acid molecules of documents cited or referenced herein, including PCT WO 99/29717, fragments thereof, e.g., ORFs and/or fragments encoding an immunogen or epitope, as well as nucleic acid molecules of strains 1103 and/or 1121, and fragments thereof, as well as vectors comprising these nucleic acid molecules, compositions comprising these nucleic molecules, vectors, or expression products therefrom, compositions comprising such expression products, primers or probes for such nucleic acid molecules, and uses or methods involving these embodiments, e.g., for detecting, diagnosing, assaying for PCV-2, for inducing an immunogenic or protective response, and the like. Indeed, this invention encompasses any inventions disclosed and/or claimed in PCT WO 99/29717 or any National application claiming priority therefrom or from the U.S. Provisionals from which that PCT claims priority.

As earlier mentioned, embodiments of the invention can include antibodies. Such antibodies can be polyclonal or monoclonal antibodies; for instance, prepared from the aforementioned circovirus, or from a polypeptide encoded by a DNA fragment having a sequence selected from the group consisting of SEQ ID NOS. 1, 2, 3, 4, 6, 7 and 8, (Figures 1-8) or from a polypeptide from expression by a vector comprising a sequence selected from the group consisting of SEQ ID NOS. 1, 2, 3, 4, 6, 7 and 8; or from a polypeptide from expression by a vector comprising DNA including an ORF selected from the group consisting of ORFs 1 to 13 (as designated in U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214). The skilled artisan may use techniques known in the art to elicit antibodies and to generate monoclonal or polyclonal antibodies. Antibodies and antigens can be used in diagnostics.

U.S. application Serial No. 09/161,092, filed September 25, 1998, U.S. application Serial No. 09/082,558, filed May 21, 1998, French applications Nos. 97 12382, 98 00873 and 98 03707, filed October 3, 1997, January 22, 1998 and March 20, 1998, respectively, and, WO-A-99 18214 also provide for probes or primers which can be useful, for instance, in detecting PCV-2 DNA, as well as for amplifying PCV-2 DNA, e.g., for preparing an expression vector. A probe or primer can be any stretch of at least 8, preferably at least 10, more preferably at least 12, 13, 14, or 15, such as at least 20, e.g., at least 23 or 25, for instance at least 27 or 30 nucleotides in PCV-2 genome or a PCV-2 gene which are unique to PCV-2 or which are in PCV-2 and are least conserved among the PCV or circovirus family. As to PCR or hybridization primers or probes and optimal lengths therefor, reference is also made to Kajimura et al., GATA 7(4):71-79 (1990). Hybridization is advantageously under conditions of high stringency, as the term "high stringency" would be understood by those with skill in the art (see, for example, Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. and Hames and Higgins, eds., 1985, Nucleic Acid Hybridization, IRL Press, Oxford, U.K.). Hybridization will be understood to be accomplished using well-established techniques, including but not limited to Southern blot hybridization, Northern blot hybridization, *in situ* hybridization and, advantageously, Southern hybridization to PCR-amplified DNA fragments.

Like probes or primers, peptides which are not full-length PCV-2 proteins are part of invention and can be any stretch of at least 8, preferably at least 10, more preferably at least 12, 13, 14, or 15, such as at least 20, e.g., at least 23 or 25, for instance at least 27 or 30 amino acids in PCV-2 which are unique to PCV-2 or which are in PCV-2 and are least conserved among the PCV and/or circovirus family. Alternatively or additionally, the amino acids of the invention which are not full length PCV-2 proteins can be an epitopic region of a PCV-2 protein.

And, as to DNA and protein sequences used in the invention, they can have homology, identity or similarity and degrees thereof as defined in U.S. application Serial No. 09/347,594, filed July 1, 1999 with homology, identity or similarity advantageously determined as discussed in USSN 09/347,594.

The PCV-2 sequences derived from Meehan et al., 1998 (Strain Imp.1010 ; ORF1 nucleotides 398-1342; ORF2 nucleotides 1381-314; and correspond respectively to ORF4 and ORF13 in U.S. application Serial No. 09/161,092 of 25 September 1998 and to COL4 and

COL13 in WO-A-9918214). Several PCV-2 strains and their sequences are disclosed herein and called Imp1008, Imp999, Imp1011-48285, Imp1011-48121, 1103 and 1121. Other strains are disclosed in A.L. Hamel *et al.* J. Virol. June 1998, vol **72**, 6: 5262-5267 (GenBank AF027217) and in I. Morozov *et al.* J. Clinical Microb. Sept. 1998 vol. **36**, 9: 2535-2541, as well as GenBank AF086834, AF086835 and AF086836. These sequences, or ORFs therefrom, or regions thereof encoding an antigen or immunogen or epitope of interest can also be used in the practice of this invention.

The invention also encompasses the equivalent sequences to those used or mentioned herein and in documents cited herein; for instance, sequences that are capable of hybridizing to the nucleotide sequence under high stringency conditions (see, e.g., Sambrook et al. (1989). Among the equivalent sequences, there may also be mentioned the gene fragments conserving the immunogenicity of the complete sequence, e.g., an epitope of interest.

The homology between the whole genome of PCV types 1 and 2 is about 75%. But within type 2, homology is generally above 95%. Thus, in the practice of the invention, use of any PCV-2 strain is encompassed by equivalence. A criteria can be that the strain is of type 2, e.g. that homology at the nucleotide level of the whole genome is equal or greater than 85%, advantageously 90% or greater, more advantageously 95% or greater, preferably 97, 98 or 99% or greater, with the strains disclosed herein, e.g. strain Imp1010.

04630238-1006000

Limits of the ORFs of strain Imp1010 are given in the following table 1:

Name	Start	End	Strand	Size of the ORF (nucleotides (nt))	Protein size (amino acids (aa))
ORF1	103	210	Sense	108 nt	35 aa
ORF2	1180	1317	Sense	138 nt	45 aa
ORF3	1363	1524	Sense	162 nt	53 aa
ORF4	398	1342	Sense	945 nt	314 aa
ORF5	900	1079	Sense	180 nt	59 aa
ORF6	1254	1334	Sense	81 nt	26 aa
ORF7	1018	704	Antisense	315 nt	104 aa
ORF8	439	311	Antisense	129 nt	42 aa
ORF9	190	101	Antisense	90 nt	29 aa
ORF10	912	733	Antisense	180 nt	59 aa
ORF11	645	565	Antisense	81 nt	26 aa
ORF12	1100	1035	Antisense	66 nt	21 aa
ORF13	314	1381	Antisense	702 nt	213 aa

The ORFs are defined with respect to strain Imp1010. The invention also encompasses the use of the corresponding ORFs in any other PCV-2 strain, and any of the PCV-2 strains as defined herein or in documents cited herein. Thus, from the genomic nucleotide sequence, it is routine art to determine the ORFs using a standard software, such as MacVector®. Also, alignment of genomes with that of strain 1010 and comparison with strain 1010 ORFs allows the one skilled in the art to readily determine the ORFs on the genome for another strain (e.g. those disclosed in WO-A-99 18214, say Imp 1008, Imp 1011-48121, Imp 1011-48285, Imp 999, as well as the new strains 1103 and 1121). Using software or making alignment is not undue experimentation and directly provides access to equivalent ORFs.

For example, referring to Figures 6 and 7, the corresponding ORFs of strains 1103 and 1121 are as given in the following table 2:

Name	Start	End	Strand	Size of the ORF (nucleotides (nt))	Protein Size (amino acids (aa))
ORF1	1524	1631	Sense	108nt	35aa
ORF2	833	970	Sense	138nt	45aa
ORF3	1016	1177	Sense	162nt	53aa
ORF4	51	995	Sense	945nt	314aa
ORF5	553	732	Sense	180nt	59aa
ORF6	907	987	Sense	81nt	26aa
ORF7	671	357	Antisense	315nt	104aa
ORF8	92	1732	Antisense	129nt	42aa
ORF9	1611	1522	Antisense	90nt	29aa
ORF10	565	386	Antisense	180nt	59aa
ORF11	298	218	Antisense	81nt	26aa
ORF12	753	688	Antisense	66nt	21aa
ORF13	1735	1037	Antisense	702nt	213aa

Also equivalent and useful in the practice of the invention are the nucleotide sequences which change neither the functionality nor the strain specificity (say of strain type 2) of the gene considered or those of the polypeptides encoded by this gene. The sequences differing through the degeneracy of the code are, of course, be included in the practice of the invention.

For ORF4, homology between PCV-1 and PCV-2 is about 86%, and for ORF13, the homology between PCV-1 and PCV-2 is about 66%. Thus, also equivalent sequences useful in the practice of the present invention, for ORF4, are those sequences having an homology equal or greater than 88%, advantageously 90% or greater, preferably 92% or 95% or greater homology with ORF4 of strain Imp1010, and for ORF13, those sequences having an homology equal or greater than 80%, advantageously 85% or greater, preferably 90% or 95% or greater than ORF13 of strain Imp1010. (Using the terminology of U.S. application Serial No. 09/161,092 of 25 September 1998.)

For homology regarding the other ORFs, one can determine those sequences which come from a PCV strain having an ORF4 and/or an ORF13 which have an homology as defined above with the corresponding ORF of strain 1010. For ORF7, sequences useful in the practice of the invention include those sequences having an homology that is advantageously equal to or greater than 80%, more advantageously 85% or greater, preferably 90% or 95% or greater with ORF7 of

strain Imp1010. For ORF10, sequences useful in the practice of the invention include those sequences having an homology that is advantageously equal to or greater than 86%, more advantageously 90% or greater, preferably 95% or greater with ORF10 of strain Imp1010. (Using the terminology of U.S. application Serial No. 09/161,092 of 25 September 1998.)

Also, equivalent sequences useful in the practice of this present invention, for ORF1 of Meehan et al., 1998, are those sequences having an homology equal or greater than 88%, advantageously 90% or greater, preferably 92% or 95% or greater with ORF1 of strain Imp1010, and for ORF2 of Meehan et al., 1998, are those sequences having an homology equal or greater than 80%, advantageously 85% or greater, preferably 90% or 95% or greater with ORF2 of strain Imp1010.

ORF1 and ORF2 according to Meehan et al., 1998 has the potential to encode proteins with predicted molecular weights of 37.7 kD and 27.8 kD respectively. ORF3 and ORF4 (according to Meehan et al. 1998, correspond to ORF7 and ORF10 respectively in WO-A-9918214 and/or U.S. application Serial No. 09/161,092 of 25 September 1998) has the potential to encode proteins with predicted molecular weights of 11.9 and 6.5 kD respectively. The sequence of these ORFs is also available in Genbank AF 055392. They can also be incorporated in plasmids and be used in accordance with the invention alone or in combination, e.g. with ORF1 and/or ORF2 of Meehan et al., 1998.

The other ORFs 1-3 and 5, 6, 8-9, 11-12 disclosed in U.S. application Serial No. 09/161,092 of 25 September 1998 (COLs 1-3 and 5, 6, 8-9, 11-12 in WO-A-9918214), or region(s) thereof encoding an antigen or epitope of interest, may be used in the practice of this invention, e.g., alone or in combination or otherwise with each other or with the ORFs 1 and/or 2 of Meehan et al., 1998 or region(s) thereof encoding antigen(s) or epitope(s). Similarly, for homology, one can determine that there are equivalent sequences which come from a PCV strain having an ORF2 and/or an ORF1 which have an homology as defined above with the corresponding ORF of strain 1010 as defined in Meehan et al., 1998. For ORF3 according to Meehan et al., 1998, an equivalent sequence has homology thereto that is advantageously, for instance, equal or greater than 80%, for example 85% or greater, preferably 90% or 95% or greater with ORF3 of strain Imp1010. And, for ORF4 according to Meehan et al., 1998, advantageously an equivalent sequence has homology that is equal or greater than 86%, advantageously 90% or greater, preferably than 95% or greater with ORF4 of strain Imp1010.

Nucleotide sequence homology can be determined using the "Align" program of Myers and Miller, ("Optimal Alignments in Linear Space", CABIOS 4, 11-17, 1988, incorporated herein by reference) and available at NCBI. Alternatively or additionally, the term "homology" or "identity", for instance, with respect to a nucleotide or amino acid sequence, can indicate a quantitative measure of homology between two sequences. The percent sequence homology can be calculated as $(N_{ref} - N_{dif}) * 100 / N_{ref}$, wherein N_{dif} is the total number of non-identical residues in the two sequences when aligned and wherein N_{ref} is the number of residues in one of the sequences. Hence, the DNA sequence AGTCAGTC will have a sequence similarity of 75% with the sequence AATCAATC ($N_{ref} = 8$; $N_{dif} = 2$).

Alternatively or additionally, "homology" or "identity" with respect to sequences can refer to the number of positions with identical nucleotides or amino acids divided by the number of nucleotides or amino acids in the shorter of the two sequences wherein alignment of the two sequences can be determined in accordance with the Wilbur and Lipman algorithm (Wilbur and Lipman, 1983 PNAS USA 80:726, incorporated herein by reference), for instance, using a window size of 20 nucleotides, a word length of 4 nucleotides, and a gap penalty of 4, and computer-assisted analysis and interpretation of the sequence data including alignment can be conveniently performed using commercially available programs (e.g., Intelligenetics™ Suite, Intelligenetics Inc. CA).. When RNA sequences are said to be similar, or have a degree of sequence identity or homology with DNA sequences, thymidine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence.

RNA sequences within the scope of the invention can be derived from DNA sequences, by thymidine (T) in the DNA sequence being considered equal to uracil (U) in RNA sequences.

Additionally or alternatively, amino acid sequence similarity or identity or homology can be determined using the BlastP program (Altschul *et al.*, Nucl. Acids Res. 25, 3389-3402, incorporated herein by reference) and available at NCBI. The following references (each incorporated herein by reference) provide algorithms for comparing the relative identity or homology of amino acid residues of two proteins, and additionally or alternatively with respect to the foregoing, the teachings in these references can be used for determining percent homology or identity: Needleman SB and Wunsch CD, "A general method applicable to the search for similarities in the amino acid sequences of two proteins," *J. Mol. Biol.* 48:444-453 (1970); Smith TF and Waterman MS, "Comparison of Bio-sequences," *Advances in Applied Mathematics*

The invention further comprehends uses of a PCV-2 immunogen, either alone or in further combination with an immunogen of another porcine pathogen to generate compositions according to the invention, e.g., admixing the ingredients; and, the invention also therefore comprehends kits wherein components are individually contained and optionally the containers are packaged together for admixture and/or administration, wherein the kit can also optionally include instructions for admixture and/or administration.

According to the present invention, immunogenic and vaccine compositions may comprise immunogens from more than one PCV-2 strain. For example, it is possible to combine immunogens from strains 1121 and 1103, from one or both of these strains with at least one other strain disclosed herein, or any other combination.

al0151

for PCV immunogens, including PCV-1 immunogens able to elicit an immunogenic or protective response against PCV-2.

Thus one aspect of the invention is to provide immunogenic or vaccinal compositions comprising a PCV immunogen and able to elicit an immunogenic or protective response against PCV-2. The invention relates also to methods of immunization or vaccination using such an immunogen, as well as to the use of such an immunogen to produce such an immunogenic or vaccinal composition.

The invention shall be further described by way of the following Example And Results, provided for illustration and not to be considered a limitation of the invention.

EXAMPLE AND RESULTS

EXAMPLE/RESULT 1: MYOCARDITIS, ABORTION AND INTRAUTERINE INFECTION ASSOCIATED WITH PCV-2

Late term abortions and farrowings with both stillborn and mummified piglets occurred in a new 450-female pig swine facility as it was brought into production. Pseudopregnancy was also observed in several gilts. Gilts received two doses of an inactivated vaccine containing parvovirus and leptospiral immunogens prior to breeding.

A litter received for postmortem examination consisted of nine fetuses that appeared to have died at various stages of gestation. There were 2 mummified, 2 macerated, 3 autolysed and 2 fresh, stillborn piglets. Lesions were observed on gross pathological examination in one partially autolysed fetus only. In this fetus both ventricles of the heart were dilated, the liver was enlarged and firm and there was both hydrothorax and ascites. Histopathologically, there were extensive areas of myocardial degeneration or necrosis with edema and mild fibrosis, and a diffuse moderate -infiltration of lymphocytes and macrophages. There was marked generalized hepatic congestion and hepatocellular loss. The spleen and kidneys were also congested. Significant histological lesions were *not* detected in the other fetuses.

Immunohistochemical staining for PCV-2 was performed as previously described using a rabbit polyclonal antiserum and a monoclonal antibody that were raised against PCV-2. on sections of formalin-fixed, routinely processed and embedded tissue (Ellis et al., 1998; Ellis et al., 1999). In the fetus with dilated cardiomyopathy there was extensive staining for PCV-2 antigen throughout the affected myocardium. Staining was most extensive in areas of necrosis and appeared to involve primarily myocytes. Both cytoplasmic and nuclear staining was present.

In multiple fetuses there was extensive staining in the liver. In some sections it appeared to involve primarily sinusoidal endothelium and Kupfer cells, while in other fetuses, including the one with myocarditis, there was also nuclear and cytoplasmic staining of hepatocytes. Positively stained cells were scattered throughout the lung, and multifocally in the kidney. Polymerase chain reaction for PCV-2 was performed as previously described using frozen tissue (Ellis et al., 1999). PCR product of the expected size for PCV-2 was amplified from fetal tissue. PCV-2 was isolated from the fetus with myocarditis and a pool of tissues from other fetuses in the litter by inoculating tissue homogenates onto PCV-free PK-15 cells.

Fetal tissues were also examined for other viral pathogens that have been associated with fetal injury and abortions in swine, including, porcine parvovirus (PPV), porcine reproductive and respiratory syndrome virus (PRRSV), encephalomyocarditis (EMCV), and enteroviruses. PPV antigen was not detected by fluorescent antibody testing (FAT) on frozen sections of lung, liver, and spleen from the mummified or stillborn fetuses. Homogenates of liver, lung, and spleen from the aborted fetuses were also inoculated into cultures of PCV-free PK-15 cells, primary porcine fallopian tube cells and Vero cells. Cytopathic viruses were not detected after three passages. Tissues were negative for PPV using PCR. PRRSV antigen was not detected by immunohistochemical staining.

Thus, there were fetal lesions and abortion directly associated with PCV-2. These results also show vertical transmission of the virus.

In a previous study, PCV-1 was isolated from 2 of 160 pig fetuses examined, implying that this group of viruses can be vertically transmitted; however, PCV-1 antigen could not be associated with any lesions in the tissue (Allan et al., 1995). The exclusion of other agents that have been associated with fetal lesions and abortion in swine, including, PPV (Bolt et al., 1997; Molitor et al., 1991), PRRSV (Lager et al., 1996), EMCV (Kim et al., 1989), and enterovirus (Molitor et al., 1991) indicate that PCV-2 can cause significant fetal pathology and subsequent abortion.

However, PCV-1 immunogens (still according to the general definition given at the beginning) may elicit an immunogenic or protective response against myocarditis and/or abortion and/or intrauterine infection as well as post-weaning multisystemic wasting syndrome and ergo PCV-1 immunogens can also be used in the practice of this invention (e.g., in the methods, compositions, uses, etc.) - either alone or in conjunction with PCV-2 immunogens (the

vector can contain and express DNA encoding for both a PCV-1 immunogen and/or epitope and a PCV-2 immunogen and/or epitope) and/or alone or in conjunction an immunogen and/or epitope of other porcine pathogen (if a vector is used, the vector can contain and express DNA encoding for both a PCV-1 immunogen and/or epitope and an immunogen and/or epitope of another porcine pathogen, or for a PCV-1 immunogen and/or epitope and a PCV-2 immunogen and/or epitope and an immunogen and/or epitope of another porcine pathogen). Thus, one skilled in the art may alternatively or additionally use a PCV-1 immunogen, and/or epitope and/or vector encoding such an immunogen and/or epitope in the practice of this invention without any undue experimentation; for instance, to so do, one need only read the text herein prior to this Example and at the conclusion of (after) this Example, and substitute --PCV-1-- for "PCV-2" with any modification minor based on teachings herein.

The wasting syndrome associated with PCV-2 infection most often occurs in 5-12 week old pigs (Allan et al., 1998; Ellis et al., 1998). Experimental infection of neonatal swine indicates a relatively long prodromal period between infection and the development of clinical signs associated with PCV-2 (Allan et al. 1999; Ellis et al. 1999). The findings herein show that the virus is transmitted vertically or in the perinatal period. Not only may interuterine vertical transmission of PCV-2 result in abortion, but it is possible that sublethally *in utero*-infected piglets may be the animals that subsequently develop PMWS.

Furthermore, these results show that inoculation of female pigs with a composition comprising an PCV-2 immunogen (which composition can also include an immunogen from another porcine pathogen, e.g., porcine parvovirus), prior to breeding or serving, or prior to the perinatal period and/or during gestation can prevent myocarditis and/or abortion and/or intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and other pathologic sequelae associated with PCV-2, by eliciting an immunological response or antibodies against PCV-2.

Of course, compositions, methods, and other aspects of the invention can be used or practiced in animals other than pigs, e.g., sheep, bison, cattle, wild boar; for instance, if PCV-2 infects such other animals.

EXAMPLE/RESULT 2: MYOCARDITIS, ABORTION AND INTRAUTERINE INFECTION ASSOCIATED WITH PCV-2

The presence of PCV-2 in neonatal piglets suggests that vertical transmission may be an important means of viral transmission. This mode of transmission may be related not only to reproductive failure, but also to the development of multisystemic disease later in life. It is of interest to determine whether previously undetected PCV-2 (and PCV-1) has been vertically transmitted in pork producing areas where PMWS, and by extension PCV-2 infection, has been endemic for at least several years.

Thirty eight submissions involving reproductive failure received in the diagnostic laboratory at the Western College of Veterinary Medicine (WCVN), University of Saskatchewan, Saskatoon, Canada, over a four-year period from a total of 30 high health herds in Canada were evaluated. Five of the farms from which the samples were obtained had diagnosed cases of PMWS. Twenty-seven of the thirty-eight submissions (71%) were classified as abortions; five of these (13%) also involved at least one mummified fetus. Of the remaining 10 cases: 5 involved stillborn piglets along, with nonviable piglets (13%); 2 with stillborn and one or more mummified feti (5%); 2 with only stillborn piglets (5%); and one with only mummified feti (2.5%). Routine diagnostics for pathogens other than circovirus revealed 4 cases (11%) in which the etiology was determined to be porcine parvovirus and 2 cases (5%) in which the etiology was determined to be of bacterial origin. Gross necropsies were performed and tissues were collected and fixed in buffered formalin (fixation time 24-72 hrs) and, in most cases, fresh tissues were also submitted for routine microbiological evaluation. None of these cases had been previously tested for PCV-2.

The PCR technique used for the detection of PCV-1 and PCV-2 was performed as previously described (Tischer et al. 1974). PCV-1 was not detected by PCR in any submissions comprising reproductive failure from the four-year period. PCV-2 was detected by PCR in two different submissions that originated from the same multi-site pork production unit on two separate occasions in the spring of the last year in the four-year period. The first of these submissions comprised a litter of piglets with gross evidence of myocarditis, cardiac hypertrophy, and chronic passive congestion.

Immunohistochemical identification of PCV-2 in tissues was performed as previously described (Tischer et al. 1974). Immunohistochemical staining (IHC) for PCV-2 was positive in

hearts from all six of the piglets that were submitted, while 4 of 6 were positive by PCV-2 PCR (see following Table).

Table: Detection of PCV-2 in the formalin fixed hearts of porcine with myocarditis by PCR, IHC and viral isolation in cell culture.

PCV-2 postive tissues	PCR	IHC	Virus Isolation
Fixed	5/6	6/6	N/A
Frozen	4/4	N/A	2/4

The second submission from the same farm consisted of a litter of four piglets in which 2 were stillborn and 2 others died shortly after birth. All four piglets also had gross evidence of a severe, difuse myocarditis, cardiac hypertrophy, and chronic passive congestion. Only fresh frozen heart, and pooled lung/spleen tissues were submitted for analysis. PCV-2 PCR was positive in the hearts of 2 of 4 piglets and in the pooled lung and splenic tissues of 4 of 4 piglets. Isolation of PCV-2 from affected hearts and/or pooled lung and splenic tissue was positive in 2 of the 4 cases that were PCV-2 positive by PCR. Based on serology and/or PCR, other agents associated with reproductive failure in swine, including porcine reproductive and respiratory syndrome virus and porcine parvovirus were apparently circulating in the breeding herd. However, these agents could not be shown to be associated with the severe cardiac (or other) lesions in the affected piglets; but, they may contribute to PMWS.

PCV-2 was not detected by PCR or IHC in any representative cases of reproductive failure submitted during the first three years of the four-year period (it was detected in cases of reproductive failure submitted during the last year of the four-year period). In order to rule out damage to DNA due to formalin fixation as a possible adverse factor limiting the ability to detect PCV-2 by PCR, PCR was performed on tissues collected from four weanling piglets with PMWS, PCV-2 DNA was amplified in all fixed tissues tested, including; lung, liver, kidney and bronchial lymph node, from all four individuals. Moreover, the sensitivity of the PCR PCV-2 was independent of the length of time that each tissue was fixed in formalin.

These results confirm and extend the previous observation (West et al. 1999) that PCV-2 can be vertically transmitted and can be present in large amounts within lesions from piglets infected *in utero*. Vertical transmission of PCV-2 virus and resultant fetal damage, such as myocarditis, is an additional disease manifestation of PCV-2. Furthermore, the failure to detect

PCV-2 in cases of reproductive failure prior to the last year of the four-year period from an endemic area of PCV-2 infection may indicate that vertical transmission was not the primary mechanism responsible for the initial dissemination of viral infection. Sexual, as well as vertical, modes of transmission can be attributed to the spread of PCV-2 infection in pigs.

EXAMPLE/RESULT 3: CULTURE AND ISOLATION OF THE PORCINE CIRCOVIRUS STRAINS

Viruses 1103 and 1021 were isolated respectively in Alberta, respectively Saskatoon, Canada, from abortive cases according to the method described in J. Ellis et al. Can. J. Vet. 1998, vol 39, 44-51.

Viral culture is carried out on PK/15 cell cultures, known to be uncontaminated with the porcine circovirus (PCV), pestiviruses, porcine adenoviruses and porcine parvoviruses (Allan G. *et al* Pathogenesis of porcine circovirus experimental infections of colostrum-deprived piglets and examination of pig foetal material. Vet. Microbiol. 1995, 44, 49-64).

Monolayers of PK/15 cells are dissociated by trypsinization (with a trypsin-versene mixture) from confluent cultures, and taken up in MEM-SA medium containing 15% foetal calf serum not contaminated by pestivirus (= MEM-G medium) in a final concentration of about 400,000 cells per ml. 10 ml aliquot fractions of this cell suspension are then mixed with 2 ml aliquot fractions of the inocula described above, and the final mixtures is aliquoted in 6 ml volumes in two Falcon flasks of 25 cm². These cultures are then incubated at +37°C for 18 hours under an atmosphere containing 10% CO₂.

After incubation, the culture medium of the semi-confluent monolayers were treated with 300 mM D-glucosamine (Cat # G48175, Sigma-Aldrich Company Limited, Poole, UK) (Tischr I. et al., Arch. Virol., 1987 96 39-57), then incubation was continued for an additional period of 48-72 hours at +37°C. Following this last incubation, one of the two Falcons of each inoculum was subjected to 3 successive freeze/thaw cycles. The PK/15 cells of the remaining Falcon were treated with a trypsin-versene solution, resuspended in 20 ml of MEM-G medium, and then inoculated into 75 cm² Falcons at a concentration of 400,000 cells/ml. The freshly inoculated flasks were then "superinfected" by addition of 5 ml of the corresponding lysate obtained after the freeze/thaw cycles.

EXAMPLE/RESULT 4: TECHNIQUE FOR THE DETECTION OF PCV BY IMMUNOFLUORESCENCE

The initial screening of all the cell culture preparations fixed with acetone was carried out by an indirect immunofluorescence technique (IIF) using a 1/100 dilution of a pool of adult pig sera. This pool of sera comprises sera from 25 adult sows from Northern Ireland and is known to contain antibodies against a wide variety of porcine viruses, including PCV: porcine parvovirus, porcine adenovirus, and PRRS virus. The IIF technique was carried out by bringing the serum (diluted in PBS) into contact with the cell cultures for one hour at +37°C, followed by two washes in PBS. The cell cultures were then stained with a 1/80 dilution in PBS of a rabbit anti-pig immunoglobulin antibody conjugated with fluorescein isothiocyanate for one hour, and then washed with PBS and mounted in glycerol buffer prior to the microscopic observation under ultraviolet light.

EXAMPLE/RESULT 5: PRODUCTION OF PCV ANTIGENS BY *IN VITRO* CULTURE

The culture of the noncontaminated PK/15 cells and the viral multiplication were carried out according to the same methods as in Example 1. The infected cells are harvested after trypsinization after 4 days of incubation at 37°C and enumerated. The next passage is inoculated with 400,000 infected cells per ml.

The various PCV-2 strains disclosed herein, e.g. strains 1103 and 1121 are so cultivated.

EXAMPLE/RESULT 6: TITRATION OF PCV-2

Titration is carried out in 96-well microplates. A suspension of PK/15 cells (150 000 cells per ml) is first introduced (100 µl per well). Then dilutions of the viral culture are done and 100 µl thereof are introduced in the wells. Incubation is done at 37°C with CO₂. After 24h, there is carried out a treatment with glucosamine half an hour at 37°C (for the conditions see example 3). The culture medium is then removed and fresh medium is introduced. Incubation is conducted 72h at 37°C. Revelation of the foci is done using an anti-PCV-2 monoclonal antibody and a FITC labelled mouse conjugate.

This method can be used to titration for preparing inactivated as well as live attenuated PCV-2.

EXAMPLE/RESULT 7: INACTIVATION OF THE VIRAL ANTIGENS

At the end of the viral culture, the infected cells are harvested and lysed using ultrasound (Branson Sonifier) or with the aid of a rotor-stator type colloid mill (UltraTurrax, IKA). The suspension is then centrifuged at 3700 g for 30 minutes. The viral suspension is inactivated with

0.1% ethyleneimine for 18 hours at +37°C or with 0.5% beta-propiolactone for 24 hours at +28°C. If the virus titre before inactivation is inadequate, the viral suspension is concentrated by ultrafiltration using a membrane with a 300 kDa cut-off (Millipore PTMK300). The inactivated viral suspension is stored at +5°C.

EXAMPLE/RESULT 8: PREPARATION OF THE VACCINE IN THE FORM OF AN EMULSION BASED ON MINERAL OIL.

The vaccine is prepared according to the following formula:

- suspension of inactivated porcine circovirus: 250 ml
- Montanide® ISA 70 (SEPPIC): 750 ml

The aqueous phase and the oily phase are sterilized separately by filtration. The emulsion is prepared by mixing and homogenizing the ingredients with the aid of a Silverson turbine emulsifier.

One vaccine dose contains about $10^{7.5}$ TCID₅₀. The volume of one vaccine dose is 0.5 ml for administration by the intradermal route, and 2 ml for administration by the intramuscular route.

This vaccine is used in a vaccination programme against the multisystemic wasting syndrome in combination with the Parvovax® vaccine.

EXAMPLE/RESULT 9: PREPARATION OF THE VACCINE IN THE FORM OF A METABOLIZABLE OIL-BASED EMULSION

The vaccine is prepared according to the following formula:

- suspension of inactivated porcine circovirus 200 ml
- Dehymuls HRE 7 (Henkel): 60 ml
- Radia 7204 (Oleofina): 740 ml

The aqueous phase and the oily phase are sterilized separately by filtration. The emulsion is prepared by mixing and homogenizing the ingredients with the aid of a Silverson turbine emulsifier.

One vaccine dose contains about $10^{7.5}$ TCID₅₀. The volume of one vaccine dose is 2 ml for administration by the intramuscular route.

This vaccine is used in a vaccination programme against the multisystemic wasting syndrome in combination with the Parvovax® vaccine.

Groups of 3 or 4 piglets, caesarian-derived day 0 are placed into isolators. The piglets are vaccinated day 2 either with (A) a plasmid comprising ORF 13 or with (B) a mixture of this plasmid and another plasmid comprising ORF 4, and with a physiological solution for the control group. Each plasmid is diluted in sterile physiological solution (NaCl 0,9%) at 250 µg/µl final concentration. A 2 ml volume is injected by intramuscular route in two points of 1 ml (1 point each side of the neck). A second injection of vaccine or placebo is administered day 14. Vaccination with DNA is well tolerated by piglets and no evidence for adverse reaction to vaccination is noted. The piglets are challenged day 21 by oronasal administration of PCV-2 viral suspension, 1 ml in each nostril. After challenge piglets are weighed once a week. Rectal temperatures are recorded on days 17, 21, 22, 24, 27, 29, 31, 34, 37, 41, 44. Day 44 fecal swabs are collected from each piglet for PCV-2 shedding. The virus is detected and quantified by quantitative PCR. Day 45 necropsies are performed and tissue samples are collected for virus isolation.

- There is no significant difference for the mean body weight gains or the mean body temperatures between groups.

- The only gross finding noted in pigs at termination is bronchial lymphadenopathy. The lesions are scored according the following criteria.

1 = mild lymph nodes enlargement, restricted to bronchial lymph nodes

2 = moderate lymph nodes enlargement, restricted to bronchial lymph nodes

3 = severe lymph nodes enlargement, extended to bronchial submandibullar prescapular and inguinal lymph nodes.

std is an abbreviation for standard deviation

Groups	Lymphadenopathy scores
--------	------------------------

mean

std

N

(A)	1.2	1.3	4
(B)	2.0	1.7	3
controls	3.0	0.0	3

N = number of piglets in each group

A reduction of the lymph node lesions is observed in 3 out 4 piglets immunized with (A) and 1 out 3 piglets immunized with (B) mixture. This difference is not significant ($p > 0.05$) due to the high value of the standard deviations (std).

- Virus load in lymph nodes tissues:

Quantitative virus re-isolation is performed on tissue homogenates prepared from bronchial and mesenteric lymph nodes.

The data presented correspond to the virus titers in tissue homogenates after transformation in \log_{10} .

Groups	PCV-2 titers				
	Bronchial LN		Mesenteric LN		N
	mean	std	mean	std	
(A)	0.9	0.8	0.9	0.8	4
(B)	0.7	0.6	0.2	0.2	3
Controls	2.0	1.1	1.8	1.1	4

Bronchial lymph nodes seem to contain the most infectious virus. A reduction of the viral load is observed in bronchial and mesenteric lymph nodes from piglets immunized with either (A) or (B) mixture. This reduction is significant ($p \leq 0.05$ for the plasmids mixture).

- Viral excretion:

Post challenge fecal swabs are assessed for shedding PCV-2 by PCR based on amplification of PCV-2 ORF 13. Each assay is performed in triplicate on 2 ml of sample. Unvaccinated

controls are negative for PCV-2 prior challenge and positive after challenge confirming the validity of the PCR assay.

Value are expressed as \log_{10} (number of molecules of PCV-2 DNA in 2 μ l sample).

<i>Groups</i>	Log₁₀ number of PCV-2 DNA molecules		
	<i>mean</i>	<i>std</i>	<i>N</i>
(A)	3.3	0.3	4
(A)	2.9	0.7	3
Controls	3.6	0.6	4

The differences between groups are not significant ($p > 0.05$).

EXAMPLE/RESULT 11: VACCINATION OF PIGLETS WITH CANARYPOX LIVE VECTOR AND RESULTS

Groups of 3 or 4 piglets, caesarian-derived day 0 are placed into isolators. Day 2 the piglets are vaccinated with 10^8 pfu of (C) a canarypox comprising ORF 13, or (D) of a canarypox comprising ORF 13 and ORF 4, or parental canarypox, in 1 ml of PBS, by intramuscular route on the side of the neck. A second injection of vaccine or placebo is administered at day 14. Vaccination with canarypox is well tolerated by piglets and no evidence for adverse reaction to vaccination is noted. The piglets are challenged day 21 by oronasal administration of a PCV-2 viral suspension, 1 ml in each nostril. Day 45 necropsies are performed and samples of tissues are collected for virus isolation.

Necropsy results :

- PMWS is characterized generally by lymphadenopathy and more rarely by hepatitis or nephritis. So the gross findings in lymph nodes are scored for each piglet in the following manner : 0 = no visible enlargement of lymph nodes ; 1 = mild lymph nodes enlargement, restricted to bronchial lymph nodes ; 2 = moderate lymph nodes enlargement, restricted to

bronchial lymph nodes ; 3 = severe lymph nodes enlargement, extended to bronchial, submandibullar prescapular and inguinal lymph nodes.

Groups	Scores
(C)	0.5
	0.0
	0.0
	1.0
mean	0.38
standard deviation	0.48
(D)	0.0
	0.5
	0.5
	1.0
mean	0.5
standard deviation	0.41
Controls	2.0
	2.5
	2.5
	2.5
mean	2.38
standard deviation	0.25

Bronchial lymphadenopathy for PCV-2 is a prominent gross finding. A significant reduction of the lymph nodes lesion in relation to control group is observed after immunization with (C) and (D) ($p \leq 0.05$).

* * *

Having thus described in detail preferred embodiments of the present invention, it is to be understood that the invention defined by the appended claims is not to be limited to particular details set forth in the above description as many apparent variations thereof are possible without departing from the spirit or scope of the present invention.

009007 82203550

References

1. Allan GM, McNeilly F, Cassady JP, et al.: 1995, Pathogenesis of porcine circovirus; experimental infections of colostrum deprived piglets and examination of pig foetal material. *Vet Microbiol* 44:49-641.
2. Allan GM, McNeilly F, Kennedy S, et al.: 1998, Isolation of porcine circovirus-like viruses from piglets with a wasting disease in the United States of America and Europe. *J Vet Diag Invest* 10:3-10.
3. Allan GM, Kennedy S, McNeilly F, et al.: 1999, Experimental reproduction of wasting disease and death by co-infection of pigs with porcine circovirus and porcine parvovirus, *J Comp Path* 121:1-11 (July 1999).
4. Bolt DM, Hani H, Muller E, and Waldvogel AS: 1997, Nonsuppurative myocarditis in piglets associated with porcine parvovirus infection. *J Comp Path* 117:107-118.
5. Ellis JA, Hassard L, Clark EG, et al.: 1998, Isolation of circovirus from lesions of piglets with postweaning multisystemic wasting syndrome. *Can Vet J* 39:44-51
6. Ellis JA, Krakowka S, Lairmore M, et al.: 1999, Reproduction of lesions of postweaning multisystemic wasting syndrome in gnotobiotic piglets. *J Vet Diag Invest* 11:3-14.
7. Kim HS, Jo HS and Bergeland ME: 1989, Serologic, virologic, and histopathologic observations of encephalomyocarditis virus infection in mummified and stillborn pigs- *J Vet Diag Invest* 1:101-104.
8. Lager KM and Halbur PG: 1996, Gross and microscopic lesions in porcine fetuses infected with porcine reproductive and respiratory syndrome virus. *J Vet Diag Invest* 8:275-282.
9. Meehan BM, McNeilly F, Todd D, et al.: 1998, Characterization of novel circovirus DNA's associated wasting disease syndromes in pigs. *J Gen Virol* 79:2171-2179.
10. Mengeling WL 1992, Porcine parvovirus. In: *Diseases of swine*, ed. Leman AD, 7th ed-, pp.299-311. Iowa State University Press, Ames, IA.
11. Molitor TW, Orveerakul K, Zhang ZZ, et al.: 1991, Polymerase chain reaction (PRC) amplification for detection of porcine parvovirus. *J Virol Meth* 32:201-211
12. Pensaert M and DeMeurichy W: 1973, A porcine enterovirus causing fetal death and mummification. Experimental infection of pregnant female pigs. *Zentralbl Veterinaarmed Beih* 11:2025.

This invention is further described by the following statements:

1. An immunological, immunogenic or vaccine composition for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

2. The composition of statement 1 for the prevention of PCV-2-caused myocarditis and/or abortion and/or intrauterine infection comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen or a polypeptide comprising an epitope of a PCV-2 immunogen or an antibody elicited by a PCV-2 immunogen or an antibody elicited by an epitope of a PCV-2 immunogen or a vector expressing a PCV-2 immunogen or a vector expressing an epitope of a PCV-2 immunogen.

3. The composition of statement 2 wherein the composition comprises a PCV-2 immunogen.

4. The composition of statement 2 wherein the composition comprises a polypeptide comprising an epitope of a PCV-2 immunogen.

5. The composition of statement 2 wherein the composition comprises an antibody elicited by a PCV-2 immunogen.

6. The composition of statement 2 wherein the composition comprises an antibody elicited by an epitope of a PCV-2 immunogen.
7. The composition of statement 2 wherein the composition comprises a vector expressing a PCV-2 immunogen.
8. The composition of statement 2 wherein the composition comprises a vector expressing an epitope of a PCV-2 immunogen.
9. The composition of statement 3 wherein the PCV-2 immunogen is a porcine circovirus.
10. The composition of statement 9 wherein the PCV-2 immunogen comprises attenuated live whole PCV-2.
11. The composition of statement 9 wherein the PCV-2 immunogen comprises inactivated PCV-2.
12. The composition of statement 3 wherein the composition is a subunit immunogenic, immunological or vaccine composition.
13. The composition of any one of statements 3, 4, 5, 6, 9, 10, 11, or 12 additionally including at least one immunogen or epitope from at least one additional pig pathogen or a vector expressing such an immunogen or epitope.
14. The composition of statement 13 wherein the composition additionally includes at least one antigen immunogen or epitope from at least one additional pig pathogen.
15. The composition of statement 13 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.
16. The composition of statement 13 wherein the at least one additional pig pathogen comprises PPV.
17. The composition of statement 7 or 8 wherein the vector comprises a DNA vector plasmid, a *E. coli*, a baculovirus, a pig herpes viruses, including Aujeszky's disease virus, a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, and a swinepox virus.
18. The composition of statement 17 wherein the vector comprises a DNA vector.
19. The composition of statement 17 wherein the vector comprises a canarypox virus.

20. The composition of statement 7 or 8 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen or epitope.

21. The composition of statement 17 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen n or epitope.

22. The composition of statement 20 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.

23. The composition of statement 21 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.

24. The composition of statement 7 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

25. The composition of statement 17 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

26. The composition of statement 24 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

27. The composition of statement 25 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

28. The composition of statement 24 wherein the vector contains and expresses ORF 4 and/or 13

29. The composition of statement 24 wherein the vector contains and expresses ORF 4 and/or 13

30. The composition of statement 3 or 4 wherein the immunogen or epitope is recombinantly produced.

31. A method for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising inducing an immunological, immunogenic or protective response against PCV-2 in a pig comprising administering to the pig a composition comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

32. The method of statement 31 for the prevention of PCV-2-caused myocarditis and/or abortion and/or intrauterine infection comprising administering a composition comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen or a polypeptide comprising an epitope of a PCV-2 immunogen or an antibody elicited by a PCV-2 immunogen or an antibody elicited by an epitope of a PCV-2 immunogen or a vector expressing a PCV-2 immunogen or a vector expressing an epitope of a PCV-2 immunogen.

33. The method of statement 32 wherein the composition comprises a PCV-2 immunogen.

34. The method of statement 32 wherein the composition comprises a polypeptide comprising an epitope of a PCV-2 immunogen.

35. The method of statement 32 wherein the composition comprises an antibody elicited by a PCV-2 immunogen.

36. The method of statement 32 wherein the composition comprises an antibody elicited by an epitope of a PCV-2 immunogen.
37. The method of statement 32 wherein the composition comprises a vector expressing a PCV-2 immunogen.
38. The method of statement 32 wherein the composition comprises a vector expressing an epitope of a PCV-2 immunogen.
39. The method of statement 33 wherein the PCV-2 immunogen is a porcine circovirus.
40. The method of statement 39 wherein the PCV-2 immunogen comprises attenuated live whole PCV-2.
41. The method of statement 39 wherein the PCV-2 immunogen comprises inactivated PCV-2.
42. The method of statement 33 wherein the composition is a subunit immunogenic, immunological or vaccine composition.
43. The method of any one of statements 33, 34, 35, 36, 39, 40, 41, or 42 wherein the composition additionally includes at least one immunogen or epitope from at least one additional pig pathogen or a vector expressing such an immunogen or epitope.
44. The method of statement 43 wherein the composition additionally includes at least one immunogen or epitope from at least one additional pig pathogen.
45. The method of statement 43 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.
46. The method of statement 43 wherein the at least one additional pig pathogen comprises PPV.
47. The method of statements 37 or 38 wherein the vector comprises a DNA vector plasmid, a *E. coli*, a baculovirus, a pig herpes viruses, including Aujeszky's disease virus, a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, and a swinepox virus.
48. The method of statement 47 wherein the vector comprises a DNA vector.
49. The method of statement 47 wherein the vector comprises a canarypox virus.

50. The method of statement 37 or 38 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen or epitope.

51. The method of statement 47 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen or epitope.

52. The method of statement 50 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.

53. The method of statement 51 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, and PPV.

54. The method of statement 37 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

55. The composition of statement 47 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

56. The method of statement 54 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

57. The method of statement 55 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

58. The method of statement 54 wherein the vector contains and expresses ORF 4 and/or 13

59. The method of statement 55 wherein the vector contains and expresses ORF 4 and/or 13

60. The method of statements 33 or 34 wherein the immunogen or epitope is recombinantly produced.

61. The method of statement 32 wherein the pig is a female pig.

62. The method of statement 61 wherein the administering is prior to breeding.

63. The method of statement 61 wherein the administering is during pregnancy.

64. The method of statement 32 wherein the pig is a male pig.

65. A method for preparing the composition of statement 1 comprising admixing the pharmaceutically or veterinarily or medically acceptable carrier and the active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 ant immunogen igen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

66. A kit for preparing the composition of statement 1 or for performing the method of statement 31 comprising in a first container the pharmaceutically or veterinarily or medically acceptable carrier and in a second container the active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope; wherein the first and second containers are optionally packaged together, and the kit optionally includes

instructions for admixture of ingredients of the composition and/or administration of the composition.

67. An isolated nucleic acid molecule comprising a sequence of the genome of 1103 strain or 1121 strain or a fragment thereof comprising an open reading frame or encoding an epitope or immunogen.

68. A vector comprising the isolated nucleic acid molecule of statement 67.

69. A PCV-2 immunogen or epitope from expression of the nucleic acid molecule of statement 67 or the vector of statement 68.

70. An immunological composition comprising the DNA molecule of statement 67 or the vector of statement 68.

71. An immunological composition comprising the immunogen or epitope of statement 69.

72. A method for inducing an immunological response comprising administering the vector of statement 68 or the nucleic acid molecule of statement 67, wherein there is in vivo expression of the nucleic acid molecule.

73. A method for inducing an immunological response comprising administering the immunological composition of statement 70, wherein there is in vivo expression of the nucleic acid molecule.

74. A method for inducing an immunological response comprising administering the immunogen or epitope of statement 69.

75. A method for inducing an immunological response comprising administering the immunological composition of statement 71.

WHAT IS CLAIMED IS:

1. An immunological, immunogenic or vaccine composition for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

2. The composition of claim 1 for the prevention of PCV-2-caused myocarditis and/or abortion and/or intrauterine infection comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen or a polypeptide comprising an epitope of a PCV-2 immunogen or an antibody elicited by a PCV-2 immunogen or an antibody elicited by an epitope of a PCV-2 immunogen or a vector expressing a PCV-2 immunogen or a vector expressing an epitope of a PCV-2 immunogen.

3. The composition of claim 2 wherein the composition comprises a PCV-2 immunogen.

4. The composition of claim 2 wherein the composition comprises a polypeptide comprising an epitope of a PCV-2 immunogen.

5. The composition of claim 2 wherein the composition comprises an antibody elicited by a PCV-2 immunogen.

6. The composition of claim 2 wherein the composition comprises an antibody elicited by an epitope of a PCV-2 immunogen.
7. The composition of claim 2 wherein the composition comprises a vector expressing a PCV-2 immunogen.
8. The composition of claim 2 wherein the composition comprises a vector expressing an epitope of a PCV-2 immunogen.
9. The composition of claim 3 wherein the PCV-2 immunogen is a porcine circovirus.
10. The composition of claim 9 wherein the PCV-2 immunogen comprises attenuated live whole PCV-2.
11. The composition of claim 9 wherein the PCV-2 immunogen comprises inactivated PCV-2.
12. The composition of claim 3 wherein the composition is a subunit immunogenic, immunological or vaccine composition.
13. The composition of claim 3 additionally including at least one immunogen or epitope from at least one additional pig pathogen or a vector expressing such an immunogen or epitope.
14. The composition of claim 13 wherein the composition additionally includes at least one antigen immunogen or epitope from at least one additional pig pathogen.
15. The composition of claim 13 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.
16. The composition of claim 13 wherein the at least one additional pig pathogen comprises PPV.
17. The composition of claim 7 wherein the vector comprises a DNA vector plasmid, a *E. coli*, a baculovirus, a pig herpes viruses, including Aujeszky's disease virus, a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, and a swinepox virus.
18. The composition of claim 17 wherein the vector comprises a DNA vector.
19. The composition of claim 17 wherein the vector comprises a canarypox virus.

31. A method for the prevention and/or treatment of porcine circovirus-2 (PCV-2)-caused myocarditis, and/or abortion and/or intrauterine infection and/or post-weaning

multisystemic wasting syndrome and/or other pathologic sequelae associated with PCV-2 comprising a inducing an immunological, immunogenic or protective response against PCV-2 in a pig comprising administering to the pig a composition comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

32. The method of claim 31 for the prevention of PCV-2-caused myocarditis and/or abortion and/or intrauterine infection comprising administering a composition comprising a pharmaceutically or veterinarily or medically acceptable carrier and an active agent comprising a PCV-2 immunogen or a polypeptide comprising an epitope of a PCV-2 immunogen or an antibody elicited by a PCV-2 immunogen or an antibody elicited by an epitope of a PCV-2 immunogen or a vector expressing a PCV-2 immunogen or a vector expressing an epitope of a PCV-2 immunogen.

33. The method of claim 32 wherein the composition comprises a PCV-2 immunogen.

34. The method of claim 32 wherein the composition comprises a polypeptide comprising an epitope of a PCV-2 immunogen.

35. The method of claim 32 wherein the composition comprises an antibody elicited by a PCV-2 immunogen.

36. The method of claim 32 wherein the composition comprises an antibody elicited by an epitope of a PCV-2 immunogen.

37. The method of claim 32 wherein the composition comprises a vector expressing a PCV-2 immunogen.

38. The method of claim 32 wherein the composition comprises a vector expressing an epitope of a PCV-2 immunogen.

39. The method of claim 33 wherein the PCV-2 immunogen is a porcine circovirus.

40. The method of claim 39 wherein the PCV-2 immunogen comprises attenuated live whole PCV-2.

41. The method of claim 39 wherein the PCV-2 immunogen comprises inactivated PCV-2.

42. The method of claim 33 wherein the composition is a subunit immunogenic, immunological or vaccine composition.

43. The method of claim 33 wherein the composition additionally includes at least one immunogen or epitope from at least one additional pig pathogen or a vector expressing such an immunogen or epitope.

44. The method of claim 43 wherein the composition additionally includes at least one immunogen or epitope from at least one additional pig pathogen.

45. The method of claim 43 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.

46. The method of claim 43 wherein the at least one additional pig pathogen comprises PPV.

47. The method of claim 37 wherein the vector comprises a DNA vector plasmid, a *E. coli*, a baculovirus, a pig herpes viruses, including Aujeszky's disease virus, a porcine adenovirus, a poxvirus, including a vaccinia virus, an avipox virus, a canarypox virus, and a swinepox virus.

48. The method of claim 47 wherein the vector comprises a DNA vector.

49. The method of claim 47 wherein the vector comprises a canarypox virus.

50. The method of claim 37 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen or epitope.

51. The method of claim 47 additionally including at least one immunogen or epitope from at least one additional pig pathogen, or a vector expressing such an immunogen or epitope, wherein the vector can also be the vector expressing the PCV-2 immunogen or epitope.

52. The method of claim 50 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, encephalomyocarditis virus, and PPV.

53. The method of claim 51 wherein the at least one additional pig pathogen is selected from the group consisting of PRRS, *Mycoplasma hyopneumoniae*, *Actinobacillus pleuropneumoniae*, *E. coli*, Atrophic Rhinitis, Pseudorabies, Hog cholera, Swine Influenza, and PPV.

54. The method of claim 37 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

55. The composition of claim 47 wherein the vector contains and expresses an ORF selected from the group consisting of ORFs 1 to 13.

56. The method of claim 54 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

57. The method of claim 55 wherein the vector contains and expresses an ORF selected from ORFs 4, 7, 10 and 13

58. The method of claim 54 wherein the vector contains and expresses ORF 4 and/or 13

59. The method of claim 55 wherein the vector contains and expresses ORF 4 and/or 13

60. The method of claim 33 wherein the immunogen or epitope is recombinantly produced.

63. The method of claim 32 wherein the pig is a female pig.

64. The method of claim 61 wherein the administering is prior to breeding.

63. The method of claim 61 wherein the administering is during pregnancy.

64. The method of claim 32 wherein the pig is a male pig.

65. A method for preparing the composition of claim 1 comprising admixing the pharmaceutically or veterinarily or medically acceptable carrier and the active agent comprising

a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 ant immunogen igen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope.

66. A kit for preparing the composition of claim 1 or for performing the method of claim 31 comprising in a first container the pharmaceutically or veterinarily or medically acceptable carrier and in a second container the active agent comprising a PCV-2 immunogen, or a polypeptide comprising an epitope of a PCV-2 immunogen, or an antibody elicited by a PCV-2 immunogen, or an antibody elicited by an epitope of a PCV-2 immunogen, or a vector expressing a PCV-2 immunogen, or a vector expressing an epitope of a PCV-2 immunogen, or a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a polypeptide comprising an epitope of a PCV-1 immunogen that binds to an antibody elicited by a PCV-2 immunogen or epitope, or an antibody elicited by a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or an antibody elicited by an epitope of a PCV-1 immunogen that binds with both a PCV-1 immunogen or epitope and a PCV-2 epitope or immunogen, or a vector expressing a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope, or a vector expressing an epitope of a PCV-1 immunogen that also binds to an antibody elicited by a PCV-2 immunogen or epitope; wherein the first and second containers are optionally packaged together, and the kit optionally includes instructions for admixture of ingredients of the composition and/or administration of the composition.

67. An isolated nucleic acid molecule comprising a sequence of the genome of 1103 strain or 1121 strain or a fragment thereof comprising an open reading frame or encoding an epitope or immunogen.

68. A vector comprising the isolated nucleic acid molecule of claim 67.

69. A PCV-2 immunogen or epitope from expression of the nucleic acid molecule of claim 67.

70. An immunological composition comprising the DNA molecule of claim 67.

71. An immunological composition comprising the immunogen or epitope of claim 69.

72. A method for inducing an immunological response comprising administering the vector of claim 68 wherein there is in vivo expression of the nucleic acid molecule.

73. A method for inducing an immunological response comprising administering the immunological composition of claim 70, wherein there is in vivo expression of the nucleic acid molecule.

74. A method for inducing an immunological response comprising administering the immunogen or epitope of claim 69.

75. A method for inducing an immunological response comprising administering the immunological composition of claim 71.

76. A PCV-2 immunogen or epitope from expression of the vector of claim 68.

77. An immunological composition comprising the immunogen or epitope of claim 76.

78. A method for inducing an immunological response comprising administering the immunogen or epitope of claim 76.

79. A method for inducing an immunological response comprising administering the immunological composition of claim 77.

80. An immunological composition comprising the vector of claim 68.

81. A method for inducing an immunological response comprising administering the immunological composition of claim 80, wherein there is in vivo expression of the nucleic acid molecule.

ABSTRACT OF THE DISCLOSURE

Porcine circovirus-2 (PCV-2) is a recently identified agent that has been associated with post-weaning multisystemic wasting syndrome (PMWS) in swine populations. The potential spectrum of disease associated with PCV-2 is expanded by evidence of vertical and sexual transmission and associated reproductive failure. PCV-2 was isolated from a litter of aborted piglets from a farm experiencing late term abortions and stillbirths. Severe, diffuse myocarditis was present in one piglet associated with extensive immunohistochemical staining for PCV-2 antigen. Variable amounts of PCV-2 antigen were also present in liver, lung and kidney of multiple fetuses. The presence of other agents that have been associated with fetal lesions and abortion in swine including porcine parvovirus, porcine reproductive respiratory syndrome virus, encephalomyocarditis virus and enterovirus could not be established. Accordingly, inoculation of female pigs with a composition including an immunogen from PCV-2 or an epitope of interest from such an immunogen or with a vector expressing such an immunogen or epitope of interest (which composition can also include an immunogen from another porcine pathogen such as porcine parvovirus or an epitope of interest therefrom or a vector expressing such an immunogen or epitope of interest, wherein the vector can co-express both the other porcine, e.g., PPV, immunogen or epitope of interest and the PCV-2 immunogen or epitope of interest, *inter alia*), e.g., prior to breeding, such as within the first five weeks of life, or prior to the perinatal period, or repeatedly over a lifetime, or during pregnancy, such as between the 6th and 8th and/or the 10th and 13th weeks of gestation, can prevent myocarditis, abortion and intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and/or other conditions associated with PCV-2. In addition, inoculation of male and/or female pigs with the aforementioned compositions can be carried out to prevent transmission of PCV-2 from male to female (or *vice versa*) during mating. Thus, the invention involves methods and compositions for preventing myocarditis, abortion and intrauterine infection associated with porcine circovirus-2, as well as post-weaning multisystemic wasting syndrome and other conditions associated with PCV-2.

Figure 1

SEQ ID No. 1

Figure 2

SEQ ID No. 2

```

aattcaacct taacctttct tattctgtag tattcaaagg gcacagagcg ggggtttgag 60
ccccctcctg ggggaagaaa gtcattaata ttgaatctca tcatgtccac cgcccaggag 120
ggcgttttga ctgtggttcg cttgacagta tatccgaagg tgcgggagag gcgggtgttg 180
aagatgccat ttttccttct ccagcggtaa cggtggcggg ggtggacgag ccaggggagg 240
cggcggagga tctggccaag atggctgcgg gggcgggtgtc ttcttctccg gtaacgcctc 300
cttggtatcg tcatatctga aaacgaaaga agtgcgctgt aagtattacc agcgacttc 360
ggcagcggca gcacctcggc agcacctcag cagcaacatg cccagcaaga agaatggaag 420
aagcggaccc caaccccata aaagggtgggt gttcactctg aataatcctt ccgaagacga 480
gcgcaagaaa atacgggatc ttccaatatc cctatttgat tattttattg ttggcgagga 540
gggtaatgag gaaggacgaa cacctcacct ccagggggttc gctaattttg tgaagaagca 600
gacttttaat aaagtgaagt ggtatttggg tgcccgtgc caccatcgaga aagcgaaagg 660
aacagatcag cagaataaag aatactgcag taaagaaggc aacttactga tggagtgtgg 720
agctcctaga tctcagggac aacggagtga cctgtctact gctgtgagta ccttggttga 780
gagcgggagt ctggtgaccg ttgcagagca gcaccctgta acgtttgtca gaaatttccg 840
cgggctggct gaacttttga aagtgcgcgg gaaaatgcag aagcgtgatt ggaagactaa 900
tgtacacgtc attgtggggc cacctgggtg tggtaaaagc aaatgggctg ctaattttgc 960
agaccggaa accacatact ggaaaccacc tagaaacaag tgggtgggatg gttaccatgg 1020
tgaagaagtg gttgttattg atgactttta tggctggctg ccctgggatg atctactgag 1080
actgtgtgat cgatatccat tgactgtaga gactaaagggt ggaactgtac cttttttggc 1140
ccgcagtatt ctgattacca gcaatcagac cccgttgga tgggtactcct caactgctgt 1200
cccagctgta gaagctcttt atcggaggat tacttccttg gtattttgga agaattgtac 1260
agaacaatcc acggaggaag ggggccagtt cgtcaccctt tcccccccat gccctgaatt 1320
tccatatgaa ataaattact gagtcttttt tatcacttcg taatggtttt tattattcat 1380
taagggttaa gtgggggggc tttaagatta aattctctga attgtacata catggttaca 1440
cggatattgt attcctgggc gtatatactg ttttcgaacg cagtgccgag gcctacgtgg 1500
tctacatttc cagtagtttg tagtctcagc cacagctgat ttcttttgtt gtttggttgg 1560
aagtaatcaa tagtggaatc taggacaggt ttgggggtaa agtagcggga gtggtaggag 1620
aagggtcggg ttatggtatg gcgggaggag tagtttacat aggggtcata ggtgagggtc 1680
gtggcctttg ttacaaagtt atcatctaga ataacagcac tggagcccac tcccctgtca 1740
ccctgggtga tcggggagca gggccag 1767

```

000001-000000

Figure 3

SEQ ID No. 3

```

aattcaacct taaccttttt tattctgtag tattcaaagg gtatagagat tttgttggtc 60
ccccctcccg ggggaacaaa gtcgtcaata ttaaattctca tcatgtccac cgcccaggag 120
ggcgttctga ctgtggtagc cttgacagta tatccgaagg tgcgggagag gcgggtgttg 180
aagatgccat ttttccttct ccaacggtag cgggtggcggg ggtgggacgag ccagggggcgg 240
cggcggagga tctggccaag atggctgcgg gggcgggtgtc ttcttctgcg gtaacgcctc 300
cttggtacg tcatagctga aaacgaaaga agtgcgctgt aagtattacc agcgcacttc 360
ggcagcggca gcacctcggc agcacctcag cagcaacatg cccagcaaga agaattggaag 420
aagcggaccc caaccacata aaagggtgggt gttcacgctg aataatcctt ccgaagacga 480
gcgcaagaaa atacgggagc tcccaatctc cctattttgat tattttattg ttggcgagga 540
gggtaattgag gaaggacgaa cacctcacct ccagggggttc gctaattttg tgaagaagca 600
aacttttaat aaagtgaagt ggtattttggg tgcccgctgc cacatcgaga aagccaaagg 660
aactgatcag cagaataaag aatattgcag taaagaaggc aacttactta ttgaatgtgg 720
agctcctcga tctcaaggac aacggagtga cctgtctact gctgtgagta ccttggttga 780
gagcgggagt ctggtgaccg ttgcagagca gcacctgtta acgtttgtca gaaatttccg 840
cgggctggct gaacttttga aagtgcgagg gaaaatgcag aagcgtgatt ggaagaccaa 900
tgtacacgtc attgtggggc cacctgggtg tggtaaaagc aaatgggctg ctaattttgc 960
agaccggaa accacatact ggaaaccacc tagaaacaag tgggtgggatg gttaccatgg 1020
tgaagaagtg gttgttattg atgactttta tggttggtg cctgtgggatg atctactgag 1080
actgtgtgat cgatatccat tgactgtaga gactaaagggt ggaactgtac cttttttggc 1140
ccgcagtatt ctgattacca gcaatcagac cccgttggaa tggtactcct caactgctgt 1200
cccagctgta gaagctctct atcggaggat tacttccttg gtattttgga agaattgtac 1260
agaacaatcc acggaggaag ggggccagtt cgtcaccttc tcccccccat gccctgaatt 1320
tccatatgaa ataaattact gagtcttttt tatcacttcg taatggtttt tattattcat 1380
ttaggggttta agtggggggg ctttaagatt aaattctctg aattgtacat acatggttac 1440
acggatattg tagtcctggc cgtatatact gttttcgaac gcagtgccga ggcctacgtg 1500
gtccacattt ctagagggtt gtagcctcag ccaaagctga ttccttttgt tatttggttg 1560
gaagtaatca atagtggagt caagaacagg tttgggtgtg aagtaacggg agtggtagga 1620
gaaggggttg gggattgtat ggcgggagga gtagtttaca tatgggtcat aggttagggc 1680
tgtggccttt gttacaaagt tatcatctag aataacagca gtggagccca ctcccctatc 1740
accctgggtg atggggggagc agggccag                                     1768

```

CCGCAGTATT CTGATTACCA GCAATCAGAC CCCGTTGGAA TGGTACTCCT CAACTGCTGT 1200

Figure 5

SEQ ID No. 5

```

aattcatatt tagcctttct aatacggtag tattggaaag gtaggggtag ggggttggtg 60
ccgcctgagg gggggaggaa ctggccgatg ttgaatttga ggtagttaac attccaagat 120
ggctgcgagt atcctccttt tatggtgagt acaaattctg tagaaaggcg ggaattgaag 180
ataccgtct ttcggcgcca tctgtaacgg tttctgaagg cggggtgtgc caaatatggt 240
cttctcggga ggatgtttcc aagatggctg cgggggcggg tccttcttct gcggtaacgc 300
ctccttgccc acgtcctcct ataaaagtga aagaagtgcg ctgctgtagt attaccagcg 360
cacttcggca gcggcagcac ctggcagcg tcagtgaaaa tgccaagcaa gaaaagcggc 420
ccgcaacccc ataagagggtg ggtgttcacc cttaataatc cttccgagga ggagaaaaac 480
aaaatacggg agcttccaat ctcccttttt gattattttg tttgcggaga ggaaggtttg 540
gaagagggtg gaactcctca cctccagggtg tttgcgaatt ttgctaagaa gcagactttt 600
aacaagggtg agtgggtattt tgggtgcccgc tgccacatcg agaaagcgaa aggaaccgac 660
cagcagaata aagaatactg cagtaaagaa ggccacatac ttatcgagtg tggagctccg 720
cggaaccagg ggaagcgcag cgacctgtct actgctgtga gtaccctttt ggagacgggg 780
tctttggtga ctgtagccga gcagttccct gtaacgtatg tgagaaattt ccgcgggctg 840
gctgaacttt tgaaagttag cggaagatg cagcagcgtg attggaagac agctgtacac 900
gtcatagtgg gccgcgccg tigtgggaag agccagtggg cccgtaattt tgctgagcct 960
agggacacct actggaagcc tagtagaaat aagtgggtggg atggatatca tggagaagaa 1020
gttggtgttt tggatgattt ttatggctgg ttacctggg atgatctact gagactgtgt 1080
gaccggtatc cattgactgt agagactaaa gggggtagctg ttcctttttt ggcccgcagt 1140
atthttgatta ccagcaatca ggccccccag gaatggtact cctcaactgc tgtcccagct 1200
gtagaagctc tctatcgag gattactact ttgcaatttt ggaagactgc tggagaacaa 1260
tccacggagg taccgaagg ccgatttgaa gcagtggacc caccctgtgc ctttttccca 1320
tataaaataa attactgagt cttttttgtt atcacatcgt aatggttttt atttttatth 1380
atthtagagg tcttttagga taaattctct gaattgtaca taaatagtca gccttaccac 1440
ataatthtg gctgtggctg cattttggag cgcatagcgg aggcctgtgt gctcgacatt 1500
ggtgtgggtg tttaaattga gccacagctg gtttctthta ttatttgggt ggaaccaatc 1560
aattgtthtg tccagctcag gtttgggggt gaagtacctg gagtggtagg taaagggtg 1620
ccttatggtg tggcgggagg agtagttaat atagggtca taggccaagt tgggtggagg 1680
ggttacaaag ttggcatcca agataacaac agtggaacca acacctctth gattagagg 1740
gatgggtct ctggggtaa
1759

```

0959238.10000

Figure 6

SEQ ID No. 6

gaattcaacc ttaacctttt ttattctgta gtattcaaag ggtataaaga ttttgttggt 60
ccccctccc gggggaacaa agtcgtcaat attaaatctc atcatgtcca ccgcccagga 120
gggcgttctg actgtggtag ccttgacagt atatccgaag gtgcgggaga rgcgggtggt 180
gaaaatgcc a ttttcccttc tccaacggtg gcggtggcgg gggtgggacma nccacgggcg 240
gcggcggaag atctggccaa gatggctgcg ggggcgggtg cttcttctgc ggtaacgcct 300
ccttggtatc gtcatactg aaaacgaaag aagtgcgtg taagtattac cagcgcactt 360
cggcagcggc agcacctcgg cagcacctca gcagcaacat gccagcaag aagaatggaa 420
gaagcggacc ccaaccacat aaaagggtggg tgttcacgct gaataatcct tccgaagacg 480
agcgcaagaa aatacgggag ctcccaatct ccctatttga ttattttatt gttggcgagg 540
agggtwtgta ggaangacga acacctcacc tccaggggtt cgctaatttt gtgaagaagc 600
aaacttttaa taaagtgaag tggatttttg gtgcccgctg ccacatcgag aaagccaaag 660
gaactgatca gcagaataaa gaattattgca gtaaagaagg caacttactt attgaatgtg 720
gagctcctcg atctcaagga caacggagtg acctgtctac tgctgtgagt acctgtgttg 780
agagcgggag tctggtgacc gttgcagagc agcacctgt aacgtttgtc agaaatttcc 840
gcgggctggc tgaacttttg aaagtgcgag ggaaaatgca gaagcgtgat tgggaagacca 900
atgtacacgt cattgtgggg ccacctgggt gtggtaaaag caaatgggct gctaattttg 960
cagaccgga aaccacatac tggaaaccac ctagaacaa gtggtgggat gggtaccatg 1020
gtgaagaagt ggttgttatt gatgactttt atggctggct gccgtgggat gatctactga 1080
gactgtgtga tcgatatcca ttgactgtag agactaaagg tggaactgta cnnnnnnngg 1140
cccgcagtat tctgattacc agcaatcaga cccggttga atggtactcc tcaactgctg 1200
tcccagctgt agaagctctc tatcggagga ttacttcctt ggtatttttg aagaatgcta 1260
cagaacaatc cacggaggaa ggggggccagt tngtcaccct ttcccccca tgccctgaat 1320
ttccatatga aataaattac tgagtccttt ttatcacttc gtaatgggtt ttattattca 1380
tttaggggtt aagtgggggg tctttaagat taaattctct gaattgtaca tacatggtta 1440
cacgatatt gtagtcctgg tcgtatatac tgttttcgaa cgcagtgccg aggcctacgt 1500
gggccacatt tctagaggtt tgtagcctca gccaaagctg attccttttg ttatttggtt 1560
ggaagtaatc aatagtggag tcaagaacag gtttggggtg gaagtaacgg gagtggtagg 1620
agaaggggtg ggggattgta tggcgggagg agtagtttac atatgggtca taggttaggg 1680
ctgtggcctt tgttacaaag ttatcatcta gaataacagc agtggagccc actcccctat 1740
caccctgggt gatgggggag cagggcca 1768

CCGCAGTAT

1841 1842 1843 1844 1845 1846 1847 1848 1849 1850
 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860
 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870
 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880
 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890
 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900
 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910
 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920
 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930
 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940
 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950
 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960
 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970
 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980
 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990
 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000
 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010
 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020
 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030
 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040
 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050
 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060
 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070
 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080
 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090
 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100
 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110
 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120
 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130
 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140
 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150
 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160
 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170
 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180
 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190
 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200
 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210
 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220
 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230
 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240
 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250
 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260
 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270
 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280
 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290
 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300
 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310
 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320
 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330
 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340
 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350
 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360
 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370
 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380
 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390
 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400
 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410
 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420
 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430
 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440
 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450
 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460
 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470
 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480
 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490
 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500
 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510
 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520
 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530
 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540
 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550
 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560
 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570
 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580
 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590
 2591 2592 2593 2594 2595 2596 2597 2598 2

accagcgcac	ttcggcagcg	gcagcacctc	ggcagcacct	cagcagcaac	atgcccagca	60
agaagaatgg	aagaagcgga	ccccaccac	ataaaaggtg	ggtgttcacg	ctgaataatc	120
cttccgaaga	cgagcgcaag	aaaatacggg	agctcccaat	ctccctattt	gattatttta	180
ttgtttggcga	ggagggtaat	gaggaaggac	gaacacctca	cctccagggg	ttcgctaatt	240
ttgtgaagaa	kcaaactttt	aataaagtga	agtggatttt	gggtgcccg	tgccacatcg	300
agaaagccaa	aggaactgat	cagcagaata	aagaatattg	cagtaaagaa	ggcaacttac	360
ttattgaatg	tggagctcct	cgatctcaag	gacaacggag	tgacctgtct	actgctgtga	420
gtaccttggt	ggagagcggg	agtctggtga	ccgttgcaga	gcagcacctc	gtaacgtttg	480
tcagaaatth	ccgcgggctg	gctgaacttt	tgaaagtga	cgggaaaatg	cagaagcgtg	540
attggaagac	caatgtacac	gtcattgtgg	ggccacctgg	gtgtggtaaa	agcaaattggg	600
ctgctaattt	tgcagaccgc	gaaaccacat	actggaaacc	acctagaaac	aagtgggtggg	660
atggttacca	tggatgaagaa	gtggttggtt	ttgatgactt	ttatggctgg	ctgccgtggg	720
atgatctact	gagactgtgt	gatcgatctc	cattgactgt	agagactaaa	ggtggaactg	780
tacctttttt	ggccccgcagt	attctgatta	ccagcaatca	gaccccgttg	gaatggtact	840
cctcaactgc	tgtcccagct	gtagaagctc	tctatcggag	gattacttcc	ttgggtatttt	900
ggaagaatgc	tacagaacaa	tccacggagg	aagggggcca	gttcgtcacc	ctttcccccc	960
catgccctga	atttccatat	gaaataaatt	actgagtcyt	ttttatcact	tcgtaatggg	1020
ttttattatt	catttagggg	ttaagtgggg	ggtctttaag	attaaattcc	ctgaattgta	1080
catacagggt	tacacggata	ttgtagtcct	ggtcgtattt	actgttttcg	aacgcagtgc	1140
cgaggcctac	gtgggtccaca	tttctagagg	tttgtagcct	cagccaaagc	tgattccttt	1200
tgttattttg	ttggaagtaa	tcaatagtgg	agtcaagaac	aggtttgggg	gtgaagtaac	1260
gggagtggtg	ggagaagggt	tgggggattg	tatggcggga	ggagtagttt	acatatgggt	1320
catatgtttg	ggctgtggcc	tttggtacaa	agttatcatc	tagaataaca	gcagtggagc	1380
ccactccctc	atcaccctgg	gtgatggggg	agcaggggca	gaattcaacc	ttaacctttc	1440
ttattctgta	gtattcaaag	ggtatagaga	ttttgttggt	ccccctccc	gggggaacaa	1500
agtcgtcaat	tttaaattctc	atcatgtcca	ccgccagga	gggcgttgtg	actgtggtac	1560
gcttgacagt	atatccgaag	gtgcggggaga	ggcgggtggt	gaagatgcca	tttttccttc	1620
tccaacggta	gcggtggcgg	gggtggacga	gccaggggcg	gcggcgagg	atctggccaa	1680
gatggctgcg	ggggcggtgt	cttcttctgc	ggtaacgcct	ccttggtat	gtcatagctg	1740
aaaacgaaag	aagtgcgctg	taagtatt				1768

Figure 8

SEQ ID No. 8

```
accagcgcac ttcggcagcg gcagcacctc ggcagcacct cagcagcaac atgcccagca 60
agaagaatgg aagaagcggg cccaaccac ataaaagggtg ggtgttcacg ctgaataatc 120
cttccgaaga cgagcgcaag aaaatacggg agctcccaat ctccctatatt gattatttta 180
ttgttggcga ggagggtaat gaggaaggac gaacacctca cctccagggg ttcgctaatt 240
ttgtgaagaa gcaaactttt aataaagtga agtggatatt ggggtgcccgc tgccacatcg 300
agaaagccaa aggaactgat cagcagaata aagaatattg cagtaaagaa ggcaacttac 360
ttattgaatg tggagctcct cgatctcaag gacaacggag tgacctgtct actgctgtga 420
gtaccttggt ggagagcggg agtctggtga ccgttgacga gcagcacctt gtaacgtttg 480
tcagaaatth ccgcgggctg gctgaacttt tgaaagtga cgggaaaatg cagaagcgtg 540
attggaagac caatgtacac gtcattgtgg ggcacactgg gtgtggtaaa agcaaaggg 600
ctgctaattt tgcagacccg gaaaccacat actggaaacc acctagaaac aagtgggtgg 660
atggttacca tgggtgaagaa gtggttggtt ttgatgactt ttatggctgg ctgccgtgg 720
atgatctact gagactgtgt gatcgatct cattgactgt agagactaaa ggtggaactg 780
tacctttttt ggcccgcagt attctgatta ccagcaatca gaccccggtg gaatgggtact 840
cctcaactgc tgtcccagct gtagaagctc tctatcggag gattacttcc ttggtatttt 900
ggaagaatgc tacagaacaa tccacggagg aagggggcca gttcgtcacc ctttcccccc 960
catgccctga atttccatat gaaataaatt actgagtcct ttttatcact tcgtaatgg 1020
ttttattatt catttagggg ttaagtgggg ggtctttaag attaaattct ctgaattgta 1080
catacatggt tacacggata ttgtagtcct ggtcgtattt actgttttcg aacgcagtgc 1140
cgaggcctac gtggtccaca tttctagagg tttgtagcct cagccaaagc tgattccttt 1200
tgttatttgg ttggaagtaa tcaatagtgg agtcaagaac aggtttgggt gtgaagtaac 1260
gggagtggta ggagaagggt tgggggattg tatggcggga ggagtagttt acatatgggt 1320
cataggttag ggctgtggcc tttggtacaa agttatcatc tagaataaca gcagtggagc 1380
ccactcccc atcacctcg gtgatggggg agcagggcca gaattcaacc ttaacctttt 1440
ttattctgta gtattcaaag ggtatagaga ttttgttgg cccccctccc gggggaacaa 1500
agtcgtcaat tttaaatctc atcatgtcca ccgcccagga gggcgttggt actgtagtac 1560
gcttgacagt atatccgaag gtgcgggaga ggcgggtgtt gaagatgcca ttttctctt 1620
tccaacggta gcggtggcgg ggggtggacga gccaggggag gcggcgagg atctggccaa 1680
gatggctgcg ggggcgggtgt cttcttctgc ggtaacgcct ccttggtatg gtcatagctg 1740
aaaacgaaag aagtgcgctg taagtatt 1768
```

CCCTTCTGTA

DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes reference to PCT International Applications)

FROMMER LAWRENCE & HAUG, LLP
File No.: 454313-2340.2

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am an original, first and joint inventor (if plural, names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention ENTITLED:

**PREVENTION OF MYOCARDITIS, ABORTION AND INTRAUTERINE INFECTION
ASSOCIATED WITH PORCINE CIRCOVIRUS-2**

the specification of which:

- ☐ is attached hereto
☒ was filed with/transmitted to USPTO on October 6, 2000 as:
☒ United States Application Serial No. _____
as the National Phase or Continuation or Continuation-in-Part of PCT
Application No. _____, filed _____,
designating the U.S., and published as _____ on _____,
☒ with amendments through DATE EVEN HEREWITH (if applicable, give details).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code § 119 (a) - (d) or § 365 (b) of any foreign application(s) for patent or inventor's certificate, or § 365 (a) of any PCT International application(s) designating at least one country other than the United State of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT International applications designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) on which priority is claimed:

Prior Foreign/PCT Application(s) [list additional applications on separate page]:

<u>Country (or PCT)</u>	<u>Application Number:</u>	<u>Filed (Day/Month/Year)</u>	<u>Priority Claimed:</u>
			<u>Yes</u> <u>No</u>
PCT	PCT/FR0002392	28/08/2000	

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

60/151,564
(Application Number)

AUGUST 31, 1999
(Filing Date)

I hereby claim the benefit under Title 35, United States Code § 120 of any United States application(s) or § 365 (c) of any PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior United States or PCT International application(s) in the manner provided by the first paragraph of Title 35, United States Code § 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Prior U.S. (or U.S.-designating PCT) Application(s) [list additional applications on separate page]:
U.S. Serial No.: Filed (Day/Month/Year) PCT Application No. Status (patented, pending, abandoned)

09/583,350

31/05/2000

PENDING

I hereby appoint William S. Frommer, Registration No. 25,506 and Thomas J. Kowalski, Registration No. 32,147, and FROMMER LAWRENCE & HAUG, LLP or their duly appointed associates, my attorneys or agents, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and to insert the Serial Number of the application in the space provided above, and specify that all communications about the application are to be directed to the following correspondence address:

Thomas J. Kowalski, Esq.
c/o FROMMER LAWRENCE &
HAUG LLP
745 Fifth Avenue
New York, NY 10151
FAX (212) 588-0500

Direct all telephone calls to: (212) 588-0800
to the attention of:
Thomas J. Kowalski

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Signature: _____ Date: _____

Signature: _____ Date: _____

Signature: _____ Date: _____

Signature: _____ Date: _____

Signature: _____ Date: _____

Signature: _____ Date: _____

hhc2091

Signature: _____ Date: _____

Full name of seventh inventor: John Harding
Residence: 43 Jubilee Drive, Humboldt Saskatchewan, SK, S0K 2A0, CA
Citizenship: Canadian

Signature: _____ Date: _____

Full name of eighth inventor: Catherine Elisabeth Charreyre
Residence: 42, Rue Ferdinand Gauthier, 69720 Saint-Laurent de Mure, FRANCE
Citizenship: French

Signature: _____ Date: _____

Full name of ninth inventor: Gilles Emile Chappuis
Residence: 3, Rue Laurent Vibert, 69006 Lyon, FRANCE
Citizenship: French

Signature: _____ Date: _____

Full name of tenth inventor: George Steve Krakowka
Residence: 2676 Summit Street, Columbus, OH 43202, USA
Citizenship: U.S.

Signature: _____ Date: _____

Full name of eleventh inventor: Jean-Christophe Francis Audonnet
Residence: 119 rue de Crequi, 69006 Lyon, FRANCE
Citizenship: French

Signature: _____ Date: _____

Full name of twelfth joint inventor: Francis McNeilly
Residence: 4 Lisleen Place, Newtownards, BT3 4NH, N. Ireland, Great Britain
Citizenship: British

Post Office Address(es) of inventors [if different from residence]:

NOTE: In order to qualify for reduced fees available to Small Entities, each inventor and any other individual or entity having rights to the invention must also sign an appropriate separate "Verified Statement (Declaration) Claiming [or Supporting a Claim by Another for] Small Entity Status" form [e.g. for Independent Inventor, Small Business Concern, Nonprofit Organization, Individual Non-Inventor].

005007 23209950